

2.7.1.37.AGC

1 Nomenclature

EC number

2.7.1.37.AGC (Protein kinases are in a state of review by the NC-IUBMB. This EC class will presumably be split up into EC 2.7.11.11, EC 2.7.11.12, EC 2.7.11.13, EC 2.7.11.14, EC 2.7.11.15, EC 2.7.11.16)

Systematic name

ATP:protein phosphotransferase

Recommended name

protein kinase (PKA, PKC, AKT, GRK, AGC-related, RSK, DBF2, etc.)

3-phosphoinositide dependent protein kinase-1 <4, 98, 102> [4, 179, 180]

AKT kinase transforming protein <178> [340]

AKT1 <113> [192]

AKT2 <180> [342]

ARK <44> [99]

Akt protein kinase <113> [187]

Akt-3 <114> [227]

Akt2 <267> [480]

Akt3 <114> [226]

Capkc1p <204> [381]

DG1 protein kinase <66> [130]

DM kinase <248> [419]

DM-kinase <229> [420]

DMPK <229> [418]

EC 2.7.1.125 (high sequence-similarity to cAMP-dependent protein kinases and protein kinase C)

EC 2.7.1.126 (high sequence-similarity to cAMP-dependent protein kinases and protein kinase C)

G protein-coupled receptor kinase 1 <184> [348]

G protein-coupled receptor kinase 2 <185> [247, 348, 349]

G protein-coupled receptor kinase GRK4 <181> [344, 345, 346]

G protein-coupled receptor kinase GRK5 <191> [362]

G protein-coupled receptor kinase GRK5 <205> [382]

G protein-coupled receptor kinase GRK5 <270> [484]

G protein-coupled receptor kinase GRK6 <206> [383]

G protein-coupled receptor kinase GRK6 <233> [428]

G protein-coupled receptor kinase GRK7 <115> [189]

GRK5 <191> [362]

GRK5 <205> [382]
 GRK6 <206> [383, 384]
 GRK7 <115> [189]
 Gprk2 <185> [349]
 ISPK-1 <216> [399]
 MT-PK <248> [449]
 NPH1 <125> [244]
 PAK-1 <272> [486]
 PDK1 <98> [179]
 PICK1 <164> [308]
 PKC Apl I <260> [471]
 PKC Apl II <261> [474]
 PKC δ II <110> [191]
 PKC δ III <145> [286]
 PKC ι <202> [377]
 PKC λ <268> [481]
 PKC ζ <122> [241]
 PKC-L <174> [336]
 PKC- δ <244> [440]
 PKC- ε <158> [302]
 PKC-zeta <137> [268]
 PKC1B <190> [361]
 PKC δ <145> [287]
 PKC ν <133> [257]
 PKD2 <116> [190]
 RAC-PK γ <273> [487]
 RAC- α serine/threonine kinase <113> [187, 192, 220, 221, 222, 223, 224]
 RAC- α serine/threonine kinase <236> [222]
 RAC- α serine/threonine kinase <179> [341]
 RAC- α serine/threonine kinase <210> [390]
 RAC- β serine/threonine protein kinase <180> [342, 343]
 RAC- β serine/threonine protein kinase <211> [390]
 RAC- β serine/threonine protein kinase <267> [480]
 RAC- γ serine/threonine protein kinase <114> [188, 225, 226, 227, 228]
 RAC- γ serine/threonine protein kinase <286> [225]
 RAC- γ serine/threonine protein kinase <273> [487]
 RAC/Akt kinase <113> [221]
 RSK3 <255> [463]
 ribosomal protein S6 kinase α 1 <274> [488]
 Rsk-1 S6 kinase <274> [488]
 S6K2 <283> [497]
 S6KII α <149> [293]
 α -PKC <164> [309]
 β -adrenergic receptor kinase <38> [90]
 β -adrenergic receptor kinase 1 <44, 47, 89> [98, 99, 100, 101, 104, 105, 167]
 β -adrenergic receptor kinase 2 <48, 49, 59> [104, 106, 119, 120]

cAMP-dependent protein kinase <1> [1]
cAMP-dependent protein kinase catalytic subunit <27, 37, 58, 61, 62> [3, 67, 68, 69, 89, 116, 117, 118, 122, 123, 124]
cAMP-dependent protein kinase type 1 <15> [30, 32, 33]
cAMP-dependent protein kinase type 2 <16> [30, 34, 35]
cAMP-dependent protein kinase type 3 <14> [29, 30, 31]
cAMP-dependent protein kinase, α -catalytic subunit <8, 52> [12, 13, 14, 15, 16, 108]
cAMP-dependent protein kinase, α -catalytic subunit <34, 45, 100> [26, 84, 85, 86, 180]
cAMP-dependent protein kinase, β -1 catalytic subunit <11> [24]
cAMP-dependent protein kinase, β -2-catalytic subunit <9> [17]
cAMP-dependent protein kinase, β -catalytic subunit <12, 13> [25, 26, 27, 28]
cAMP-dependent protein kinase, β -catalytic subunit <41> [93]
cAMP-dependent protein kinase, γ -catalytic subunit <40> [93, 94]
cAPK <58> [116]
cGK <73> [140]
cGK II <88> [166]
cGMP-dependent protein kinase 1, α isozyme <6, 7, 73> [6, 7, 8, 9, 10, 11, 77, 140]
cGMP-dependent protein kinase 1, β isozyme <30, 36, 97> [11, 77, 78, 88, 178]
cGMP-dependent protein kinase 2 <70, 84, 88> [133, 134, 135, 161, 166]
cGMP-dependent protein kinase, isozyme 1 <66> [67, 115, 130]
cGMP-dependent protein kinase, isozyme 2 forms cD5/T2 <57> [115]
calcium dependent protein kinase C
calcium-dependent protein kinase C <260> [471, 472, 473]
calcium-independent protein kinase C <261> [471, 472, 473, 474, 475]
calcium/phospholipid dependent protein kinase
cell cycle protein kinase DBF2 <167> [316, 317]
guanosine cyclic 3',5'-phosphate dependent protein kinase <7> [10]
heart muscle protein kinase
insulin-stimulated protein kinase <216> [399]
myotonic dystrophy protein kinase <229> [418, 419, 420, 421]
myotonin-protein kinase <229> [418]
myotonin-protein kinase <248> [419, 421, 449, 450, 451, 452, 453, 454]
nPKC eta <170> [326]
nPKC theta <238> [433]
nonphototropic hypocotyl protein 1 <125> [243, 244, 245, 246]
p54 S6 kinase 2 <283> [497]
p70 S6 kinase <165> [311]
p70 S6 kinase <283> [499]
p70(S6k) <165> [311]
phosphatidylserine-sensitive calcium-dependent protein kinase
phototropin <125> [244]
pp90rsk Ser/Thr kinase <255> [463]
protein kinase 2 <175> [337]

protein kinase A
 protein kinase B α <113> [221]
 protein kinase B γ <286> [225, 228]
 protein kinase B γ <114> [228]
 protein kinase C
 protein kinase C <156> [247, 299]
 protein kinase C α <160> [305]
 protein kinase C δ <110> [191]
 protein kinase C δ <145> [286]
 protein kinase C, D2 type <116> [190, 229]
 protein kinase C, α type <135> [263, 264]
 protein kinase C, α type <139> [263, 270, 271]
 protein kinase C, α type <148> [263, 281]
 protein kinase C, α type <160> [304, 305]
 protein kinase C, α type <164> [308, 309, 310]
 protein kinase C, β type <109> [183, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208]
 protein kinase C, β type <136> [263, 265]
 protein kinase C, β type <141, 142> [275, 276, 277, 278, 279, 280, 281]
 protein kinase C, brain isoenzyme <138> [247, 269]
 protein kinase C, δ type <110> [184, 191, 209, 210, 211, 212]
 protein kinase C, δ type <145> [286, 287, 288, 289, 290, 291]
 protein kinase C, δ type <244> [440, 441]
 protein kinase C, ε type <146> [290, 291]
 protein kinase C, ε type <152> [295]
 protein kinase C, ε type <158> [301, 302]
 protein kinase C, ε type <239> [434]
 protein kinase C, η type <170> [326]
 protein kinase C, η type <174> [334, 335, 336]
 protein kinase C, η type <275> [489]
 protein kinase C, γ type <140> [208, 270, 272, 273, 274]
 protein kinase C, γ type <137> [265, 266, 267, 268]
 protein kinase C, γ type <151> [294]
 protein kinase C, ι type <202> [376, 377]
 protein kinase C, ι type <268> [481]
 protein kinase C, μ type <254> [462]
 protein kinase C, μ type <269> [482, 483]
 protein kinase C, ν type <133> [257]
 protein kinase C, ζ type <122> [241]
 protein kinase C, ζ type <243> [268, 439]
 protein kinase C, ζ type <147> [290, 291, 292]
 protein kinase C, ζ type <241> [436]
 protein kinase C- ε <239> [434]
 protein kinase C- η <275> [489]
 protein kinase C-like <234> [429]

protein kinase C-like <278> [429]
 protein kinase C-like 1 <193> [182, 364]
 protein kinase C-like 1 <173> [331, 332, 333]
 protein kinase C-like 1 <258> [334, 469, 470]
 protein kinase C-like 1 <189> [359, 360]
 protein kinase C-like 1 <204> [381]
 protein kinase C-like 1 <272> [470, 486]
 protein kinase C-like 2 <194> [182, 364, 365, 366]
 protein kinase C-like 2 <259> [334, 469]
 protein kinase C-like 2 <190> [361]
 protein kinase C δ <110> [184]
 protein kinase C δ <145> [287]
 protein kinase C μ <269> [482]
 protein kinase D <269> [483]
 protein kinase D2 <116> [190]
 protein kinase DBF20 <182> [316, 321]
 protein kinase DC1 <33> [69]
 protein kinase DC2 <159> [303]
 protein kinase G
 protein kinase HMK
 protein kinase PKX1 <217> [401]
 protein kinase PVPK-1 <157> [300]
 protein kinase PrkC1
 protein kinase SGK <111> [185]
 protein kinase Sgk <107> [181]
 protein kinase Ulc1
 protein kinase x
 rac-PK <180> [343]
 rhodopsin kinase <53, 76, 86, 96> [109, 110, 146, 147, 148, 165, 177]
 ribosomal S6 kinase (Rsk-2) <216> [396]
 ribosomal protein S6 kinase <165> [311, 312, 313, 314]
 ribosomal protein S6 kinase <171> [327]
 ribosomal protein S6 kinase II α <149> [293]
 ribosomal protein S6 kinase II α <161> [306]
 ribosomal protein S6 kinase II β <150> [293]
 ribosomal protein S6 kinase α 1 <162> [306]
 ribosomal protein S6 kinase α 1 <256> [400]
 ribosomal protein S6 kinase α 2 <255> [400, 463]
 ribosomal protein S6 kinase α 2 <287> [501]
 ribosomal protein S6 kinase α 3 <216> [396, 397, 398, 399, 400]
 ribosomal protein S6 kinase α 6 <285> [500]
 ribosomal protein S6 kinase β 2 <283> [497, 498, 499]
 ribosomal protein S6 kinase β 2 <290> [504]
 serine/threonine-protein kinase AtPK1/AtPK6 <203> [243, 378, 379, 380]
 serine/threonine-protein kinase AtPK19 <265> [243, 378]
 serine/threonine-protein kinase Pk61C <95> [67, 176]
 serine/threonine-protein kinase SCH9 <23> [54, 55, 56]

serine/threonine-protein kinase Sgk <107> [181, 194, 195, 196]
 serine/threonine-protein kinase Sgk <111> [185, 213, 214, 215, 216, 217, 218]
 serine/threonine-protein kinase Sgk <288> [195]
 serine/threonine-protein kinase Sgk <245> [196, 442, 443, 444, 445]
 serine/threonine-protein kinase YPK1 <153> [296, 297]
 serine/threonine-protein kinase YPK2/YKR2 <163> [296, 307]
 serine/threonine-protein kinase cot-1 <198> [370]
 serine/threonine-protein kinase nrc-2 <124> [242]
 serine/threonine-protein kinase orb6 <2> [2]
 serine/threonine-protein kinase sck1 <63> [3, 125]
 spermatozoon associated protein kinase <166> [315]
 type I β isozyme of cGMP-dependent protein kinase <30> [78]

CAS registry number

141436-78-4 (calcium dependent protein kinase C)
 141588-27-4 (cGMP-dependent protein kinase)
 142008-29-5 (cAMP-dependent protein kinase)
 191808-15-8 (phosphoinositide dependent protein kinase 1)
 377752-08-4 (ribosomal protein S6 kinase 2)
 389133-24-8 (ribosomal S6 kinase 3)
 90698-26-3 (ribosomal protein S6 kinase 1)

2 Source Organism

<1> *Paramecium primaurelia* [1]
 <2> *Schizosaccharomyces pombe* [2]
 <4> *Rattus norvegicus* [4]
 <6> *Oryctolagus cuniculus* [6]
 <7> *Bos taurus* [7, 8, 9, 10, 11]
 <8> *Bos taurus* [12, 13, 14, 15, 16]
 <9> *Bos taurus* [17]
 <11> *Bos taurus* [24]
 <12> *Mus musculus* [25, 26, 27, 28]
 <13> *Cricetulus griseus* [26]
 <14> *Saccharomyces cerevisiae* [29, 30, 31]
 <15> *Saccharomyces cerevisiae* [30, 32, 33]
 <16> *Saccharomyces cerevisiae* [30, 34, 35]
 <23> *Saccharomyces cerevisiae* [54, 55, 56]
 <27> *Drosophila melanogaster* [67, 68, 69]
 <30> *Homo sapiens* [77, 78]
 <33> *Drosophila melanogaster* [69]
 <34> *Homo sapiens* [84, 85, 86]
 <36> *Bos taurus* [11, 88]
 <37> *Caenorhabditis elegans* [89]
 <38> *Bos taurus* [90]
 <40> *Homo sapiens* [93, 94]

- <41> *Homo sapiens* [93]
- <44> *Homo sapiens* [98, 99, 100, 101]
- <45> *Cricetulus griseus* [26]
- <47> *Rattus norvegicus* [104, 105]
- <48> *Bos taurus* [106]
- <49> *Rattus norvegicus* [104]
- <52> *Rattus norvegicus* [108]
- <53> *Bos taurus* [109, 110]
- <57> *Drosophila melanogaster* [115]
- <58> *Dictyostelium discoideum* [116, 117, 118]
- <59> *Homo sapiens* [119, 120]
- <61> *Schizosaccharomyces pombe* [3, 122, 123]
- <62> *Ascaris suum* [124]
- <63> *Schizosaccharomyces pombe* [3, 125]
- <66> *Drosophila melanogaster* [67, 115, 130]
- <67> *Drosophila melanogaster* [115]
- <70> *Homo sapiens* [133, 134, 135]
- <73> *Homo sapiens* [77, 140]
- <76> *Homo sapiens* [146, 147, 148]
- <84> *Mus musculus* [161]
- <86> *Rattus norvegicus* [148, 165]
- <88> *Rattus norvegicus* [166]
- <89> *Mesocricetus auratus* [167]
- <95> *Drosophila melanogaster* [67, 176]
- <96> *Mus musculus* [177]
- <97> *Mus musculus* [178]
- <98> *Mus musculus* [179]
- <100> *Mus musculus* [180]
- <102> *Homo sapiens* [180]
- <105> *Schizosaccharomyces pombe* [180]
- <107> *Mus musculus* [181, 194, 195, 196]
- <109> *Mus musculus* [183, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208]
- <110> *Mus musculus* [184, 191, 209, 210, 211, 212]
- <111> *Homo sapiens* [185, 213, 214, 215, 216, 217, 218]
- <113> *Homo sapiens* [187, 192, 220, 221, 222, 223, 224]
- <114> *Homo sapiens* [188, 225, 226, 227, 228]
- <115> *Homo sapiens* [189]
- <116> *Homo sapiens* [190, 229]
- <122> *Oryctolagus cuniculus* [241]
- <124> *Neurospora crassa* [242]
- <125> *Arabidopsis thaliana* [243, 244, 245, 246]
- <133> *Homo sapiens* [257]
- <135> *Bos taurus* [263, 264]
- <136> *Bos taurus* [263, 265]
- <137> *Homo sapiens* [265, 266, 267, 268]
- <138> *Drosophila melanogaster* [247, 269]
- <139> *Rattus norvegicus* [263, 270, 271]

- <140> *Mus musculus* [208, 270, 272, 273, 274]
- <141> *Homo sapiens* [275, 276, 277, 278, 279, 280]
- <142> *Oryctolagus cuniculus* [281]
- <145> *Rattus norvegicus* [286, 287, 288, 289, 290, 291]
- <146> *Rattus norvegicus* [290, 291]
- <147> *Rattus norvegicus* [290, 291, 292]
- <148> *Oryctolagus cuniculus* [263, 281]
- <149> *Xenopus laevis* [293]
- <150> *Xenopus laevis* [293]
- <151> *Oryctolagus cuniculus* [294]
- <152> *Oryctolagus cuniculus* [295]
- <153> *Saccharomyces cerevisiae* [296, 297]
- <156> *Drosophila melanogaster* [247, 299]
- <157> *Phaseolus vulgaris* [300]
- <158> *Mus musculus* [301, 302]
- <159> *Drosophila melanogaster* [303]
- <160> *Homo sapiens* [304, 305]
- <161> *Gallus gallus* [306]
- <162> *Mus musculus* [306]
- <163> *Saccharomyces cerevisiae* [296, 307]
- <164> *Mus musculus* [308, 309, 310]
- <165> *Rattus norvegicus* [311, 312, 313, 314]
- <166> *Aplysia californica* [315]
- <167> *Saccharomyces cerevisiae* [316, 317]
- <170> *Mus musculus* [326]
- <171> *Homo sapiens* [327]
- <173> *Saccharomyces cerevisiae* [331, 332, 333]
- <174> *Homo sapiens* [334, 335, 336]
- <175> *Dictyostelium discoideum* [337]
- <178> *AKT8 murine leukemia virus* [340]
- <179> *Mus musculus* [341]
- <180> *Homo sapiens* [342, 343]
- <181> *Homo sapiens* (4 isoforms: GRK4 α , GRK4 β , GRK4 γ , GRK4 δ [344]; 2 isoforms: GRK4A and GRK4B [345]) [344, 345, 346]
- <182> *Saccharomyces cerevisiae* [316, 321]
- <184> *Drosophila melanogaster* [348]
- <185> *Drosophila melanogaster* [247, 348, 349]
- <189> *Caenorhabditis elegans* [359, 360]
- <190> *Caenorhabditis elegans* [361]
- <191> *Homo sapiens* [362]
- <193> *Schizosaccharomyces pombe* [182, 364]
- <194> *Schizosaccharomyces pombe* [182, 364, 365, 366]
- <198> *Neurospora crassa* [370]
- <202> *Homo sapiens* [376, 377]
- <203> *Arabidopsis thaliana* [243, 378, 379, 380]
- <204> *Candida albicans* [381]
- <205> *Bos taurus* [382]

- <206> *Homo sapiens* [383, 384]
- <210> *Rattus norvegicus* [390]
- <211> *Rattus norvegicus* [390]
- <216> *Homo sapiens* [396, 397, 398, 399, 400]
- <217> *Homo sapiens* [401]
- <229> *Mus musculus* [418, 419, 420, 421]
- <233> *Rattus norvegicus* [428]
- <234> *Aspergillus niger* [429]
- <236> *Bos taurus* [222, 224]
- <238> *Mus musculus* [433]
- <239> *Homo sapiens* [434]
- <241> *Mus musculus* [436]
- <242> *Homo sapiens* [437, 438]
- <243> *Homo sapiens* [268, 439]
- <244> *Homo sapiens* [440, 441]
- <245> *Rattus norvegicus* [196, 442, 443, 444, 445]
- <248> *Homo sapiens* [419, 421, 449, 450, 451, 452, 453, 454]
- <254> *Homo sapiens* [462]
- <255> *Homo sapiens* [400, 463]
- <256> *Homo sapiens* [400]
- <258> *Homo sapiens* [334, 469, 470]
- <259> *Homo sapiens* [334, 469]
- <260> *Aplysia californica* [471, 472, 473]
- <261> *Aplysia californica* [471, 472, 473, 474, 475]
- <265> *Arabidopsis thaliana* [243]
- <267> *Mus musculus* [480]
- <268> *Mus musculus* [481]
- <269> *Mus musculus* [482, 483]
- <270> *Rattus norvegicus* [484]
- <272> *Rattus norvegicus* [470, 486]
- <273> *Rattus norvegicus* [487]
- <274> *Rattus norvegicus* [488]
- <275> *Rattus norvegicus* [489]
- <278> *Trichoderma reesei* [429]
- <283> *Homo sapiens* [497, 498, 499]
- <285> *Homo sapiens* [500]
- <286> *Mus musculus* [225, 228]
- <287> *Mus musculus* [501]
- <288> *Oryctolagus cuniculus* [195]
- <290> *Mus musculus* [504]

3 Reaction and Specificity

Catalyzed reaction

ATP + a protein = ADP + a phosphoprotein

Reaction type

phospho group transfer

Natural substrates and products

- S** ATP + S6 protein of the 40s ribosomal subunit <165> (<165>, the mitogen-activated protein kinase plays a central role in the control of mRNA translation. It physiologically phosphorylates the S6 protein of the 40s ribosomal subunit in response to mitogenic stimuli and is a downstream component of the rapamycin-sensitive pathway, which includes the 12-kDa FK506 binding protein and includes rapamycin and the 12-kDa FK506 binding protein target 1 [311]) (Reversibility: ? <165> [311]) [311]
- P** ADP + phosphorylated S6 protein of the 40s ribosomal subunit
- S** ATP + protein <44, 48> (<44>, the enzyme mediates agonist-dependent phosphorylation of the β 2-adrenergic and related G protein-coupled receptors [98]; <44>, specifically phosphorylates the agonist-occupied forms of the β 2-adrenergic receptor and related G protein-coupled receptors [101]; <48>, specifically phosphorylates the agonist-occupied form of the β -adrenergic and related G protein-coupled receptors [106]) (Reversibility: ? <44, 48> [98, 101, 106]) [98, 101, 106]
- P** ATP + phosphoprotein
- S** ATP + protein kinase C ζ <98> (<98>, phosphorylation and activation of protein kinase C ζ [179]) (Reversibility: ? <98> [179]) [179]
- P** ADP + phosphorylated protein kinase Czeta
- S** ATP + rhodopsin <53, 76, 86, 96> (<53>, light-dependent deactivation of rhodopsin involves receptor phosphorylation that is mediated by the highly specific protein kinases rhodopsin kinase [109]; <76>, null mutations in the rhodopsin kinase gene are a cause of Oguchi disease and extend the known genetic heterogeneity in congenital stationary night blindness [147]; <96>, enzyme is required for normal rhodopsin deactivation. Abnormal photoresponses and light-induced apoptosis in rods lacking rhodopsin kinase [177]) (Reversibility: ? <53, 76, 86, 96> [109, 146, 147, 148, 177]) [109, 146, 147, 148, 177]
- P** ADP + phosphorylated rhodopsin
- S** Additional information <107, 109, 110, 111, 113, 115, 122, 124, 125, 135, 137, 145> (<107>, induction of enzymatically active Sgk functions as a key cell survival component in response to different environmental stress stimuli [181]; <109>, protein kinase C β controls nuclear factor κ B activation in B cells through selective regulation of the κ B kinase α [183]; <110>, protein kinase C δ controls self-antigen-induced B-cell tolerance [184]; <111>, K⁺-channel activation by all three isoforms of serum-dependent and glucocorticoid-dependent protein kinase SGK [185]; <115>, kinase plays a role in human visual signaling [189]; <110>, key regulatory role in a variety of cellular functions, including apoptosis, as well as cell growth and differentiation [191]; <113>, AKT1 gene is not a major contributor to susceptibility to type II diabetes mellitus in Ashkenazi Jews [192]; <107>, induction of sgk by aldosterone is detected in kidney cortex and medulla, whereas the papilla express a constitutively high level of the

kinase [194]; <107>, enzyme plays an important role in the early phase of aldosterone-stimulated Na^+ transport [195]; <107>, transcriptional target of p53 in mammary epithelial cells, hormone-regulated protein kinase gene with a functionally defined p53 promoter recognition element [196]; <109>, overproduction of protein kinase C causes disordered growth control in rat fibroblasts, activation of PKC may be of central importance in the process of multistage carcinogenesis [204]; <110>, PKC δ is involved in fundamental cellular functions regulated by diacylglycerols and mimicked by phorbol esters [211]; <111>, the enzyme acts in concert with Akt to propagate the effects of PI3K activation within the nucleus and to mediate the biological outputs of PI3K signaling, including cell survival and cell cycle progression [213]; <111>, deranged transcriptional regulation of cell-volume-sensitive kinase hSGK in diabetic nephropathy [214]; <111>, transcript levels are strongly altered during anisotonic and isotonic cell volume changes [218]; <122>, PKC ζ action is involved in growth and differentiation of the collecting duct [241]; <124>, the enzyme is required to repress entry into the conidiation program [242]; <125>, NPH1 is an autophosphorylating flavoprotein photoreceptor mediating phototropic responses in higher plants [245]; <135>, the enzyme is a phorbol ester receptor [264]; <137>, PKC ζ exhibits a constitutive kinase [268]; <145>, PKC δ III may show a dominant negative effect against PKC δ I [286]) [181, 183, 184, 185, 189, 192, 194, 195, 196, 204, 211, 213, 214, 241, 242, 245, 264, 268, 286]

P ?

S Additional information <153, 158, 163, 164, 166, 167, 173, 178, 180, 181, 182, 185, 198, 203> (<153,163>, enzyme is required for cell growth [296]; <158>, protein kinase C- ϵ increases growth and cause malignant transformation when overexpressed in NIH3T3 cells the catalytic domain of PKC- ϵ , in reciprocal PKC- δ and PKC- ϵ chimeras, is responsible for conferring tumorigenicity to NIH3T3 cells, whereas both regulatory and catalytic domains of PKC- ϵ contribute to in vitro transformation [301]; <164>, plays a central role in the control of proliferation and differentiation of a wide range of cell types by mediating the signal transduction response to hormones and growth factors [308]; <166>, kinase plays a part in regulating events associated with fertilization [315]; <167>, DBF2 mRNA is expressed under cell-cycle control at or near START [316]; <182>, the DBF20 mRNA is expressed at a low level and not under cell-cycle control [316]; <167>, DBF2 is likely to encode a protein kinase that may function in initiation of DNA synthesis and also in late nuclear division [317]; <173>, PKC1-depleted cells arrested growth with small buds. PKC₁ may regulate a previously unrecognized checkpoint in the cell cycle [333]; <178>, enzyme may form a functional link between tyrosine and serine-threonine phosphorylation pathways [340]; <180>, AKT2 may contribute to the pathogenesis of ovarian carcinomas [342]; <181>, enzyme is involved in fertilization [344]; <185>, Gprk2 is required for egg morphogenesis [349]; <203>, enzyme may function in the adaptation of plant cells to cold or high-salt conditions [378]; <203>, atpk1 is involved in

the control of plant growth and development [380]) [296, 301, 308, 315, 316, 317, 333, 340, 342, 344, 349, 370, 378, 380]

P ?

S Additional information <203, 216, 242, 243, 244, 245, 248, 254, 267, 270, 285, 290> (<216>, mutations in the ribosomal S6 kinase (Rsk-2) gene are associated with Coffin-Lowry syndrome, an X-linked disorder characterized by facial dysmorphism, digit abnormalities and severe psychomotor retardation [396,397,398]; <216> inherited defects in insulin-stimulated activation of muscle glycogen synthesis in patients with insulin-resistant NIDDM may be located further upstream of ISPK-1 in the insulin action cascade [399]; <242>, enzyme may play a role in signal transduction and growth regulatory pathways unique to hematopoietic cells [438]; <243>, constitutive enzyme [268]; <244>, PKC- δ desensitizes the Pkc1-mediated pathway by regulating an aspect of G protein function [440]; <245>, SGK is a component of the phosphoinositide 3 (PI 3)-kinase signaling pathway [442]; <245>, the enzyme is transcriptionally regulated by serum and glucocorticoids in mammary epithelial cells, hormone-regulated protein kinase gene with a functionally defined p53 promoter recognition element [196]; <245>, enzyme is induced during ovarian cell differentiation [442]; <245>, induction of *sgk* gene may be associated with a series of axonal regenerations after brain injury, and in addition, the *sgk* gene may also play important roles in the development of particular groups of neurons in the postnatal brain [444]; <245>, the enzyme is highly regulated at the transcriptional level by glucocorticoid hormones [445]; <248>, T-PK may have specialized functions in different areas of central nervous system. Alterations of this complex expression pattern can be responsible for the mental status impairment observed in myotonic dystrophy patients [449]; <248>, enzyme may have a role in the development of mental symptoms in severe cases of myotonic dystrophy [451]; <248>, decreased expression of myotonin-protein kinase messenger RNA and protein in adult form of myotonic dystrophy [452]; <254>, in normal rat kidney cells, predominant phosphorylation of a 30000 Da protein at serine residues, constitutive low level expression in normal tissues, elevated expression levels in selected tumor cell lines, a role of PKC μ in signal transduction pathways related to growth control [462]; <203>, may function in the adaptation of plant cells to cold or high-salt conditions [378]; <267>, Akt2 expression is activated during cellular differentiation and may function in the signaling pathways of some adult tissues [480]; <270>, enzyme mainly involved in homologous desensitization of the TSH receptor [484]; <285>, ribosomal S6-kinase RSK4 is commonly deleted in patients with complex X-linked mental retardation, RSK4 plays a role in normal neuronal development. RSK4 is completely deleted in eight patients with the contiguous gene syndrome including MRX, partially deleted in a patient with DFN3 and present in patients with an Xq21 deletion and normal intellectual abilities [500]; <290>, enzyme is important for cell growth [504]) [268, 378, 396, 397, 398, 399, 438, 440, 442, 444, 445, 449, 451, 452, 462, 480, 484, 500, 504]

P ?

S Additional information <2, 7, 23, 30, 34, 38, 44, 47, 49, 58, 70, 73> (<44>, role for β ARK in modulating some receptor-mediated immune functions [99]; <7>, amino acid sequence at the ATP-binding site of cGMP-dependent protein kinase [7]; <23>, the enzyme positively regulates the progression of yeast cells through the G1 phase of the cell cycle [54]; <23>, the enzyme is part of a growth control pathway which is at least partially redundant with the cAMP pathway [56]; <30,73>, enzyme plays a crucial role in the relaxation of vascular smooth muscle by lowering the intracellular level of calcium [77]; <34>, the enzyme plays a central role in the control of mammalian sperm capacitation and motility [84]; <38>, enzyme is important in mediating rapid agonist-specific desensitization [90]; <47,49>, general role in the desensitization of synaptic receptors [104]; <58>, enzyme plays an essential role during differentiation and fruit morphogenesis in *Dictyostelium discoideum* [116]; <70>, the enzyme is thought to be involved in the regulation of intestinal ion transport and fluid secretion [134]; <70>, plays a pivotal role in the regulation of intestinal fluid balance in man [135]; <73>, enzyme is involved in inhibition of platelet aggregation, relaxation of smooth muscle cells, and control of cardiocyte contractility. Pathophysiological implication of the type I cGK in cardiovascular diseases, including hypertension and atherosclerosis [140]) [2, 7, 54, 56, 77, 84, 90, 99, 104, 116, 134, 135, 140]

P ?

Substrates and products

S ATP + PKC- α -derived peptide <137> (<137>, in the presence of the classical PKC activators phosphatidylserine/diacylglycerol, PKC α phosphorylates a PKC- α pseudosubstrate-derived peptide, an epidermal-growth-factor-receptor-derived peptide, histone III-S and myelin basic protein to an equal extent, whilst PKC ζ phosphorylates only the PKC- α -derived peptide [268]) (Reversibility: ? <137> [268]) [268]

P ADP + phosphorylated PKC- α -derived peptide

S ATP + S6 protein of the 40s ribosomal subunit <165> (<165>, the mitogen-activated protein kinase plays a central role in the control of mRNA translation. It physiologically phosphorylates the S6 protein of the 40s ribosomal subunit in response to mitogenic stimuli and is a downstream component of the rapamycin-sensitive pathway, which includes the 12-kDa FK506 binding protein and includes rapamycin and the 12-kDa FK506 binding protein target 1 [311]) (Reversibility: ? <165> [311]) [311]

P ADP + phosphorylated S6 protein of the 40s ribosomal subunit

S ATP + activated form of G protein-coupled receptors <181> [346]

P ADP + phosphorylated G protein-coupled receptors

S ATP + β 2-adrenergic receptor <205, 206> (<205>, phosphorylation in an agonist-dependent manner, phosphorylates the C-terminal tail regions of both receptor proteins [382]) (Reversibility: ? <205,206> [382,384]) [382, 384]

P ADP + phosphorylated β 2-adrenergic receptor

- S** ATP + histone H1 <113, 116> (<116>, PKD2 activated by phorbol esters efficiently phosphorylate the exogenous substrate histone H1 [190]) (Reversibility: ? <113,116> [190,224]) [190, 224]
- P** ADP + phosphorylated histone H1
- S** ATP + histone H1S <275> (<275>, poor substrate [489]) (Reversibility: ? <275> [489]) [489]
- P** ADP + phosphorylated histone H1S
- S** ATP + myelin basic protein <202, 204> (Reversibility: ? <202,204> [377,381]) [377, 381]
- P** ADP + phosphorylated myelin basic protein
- S** ATP + plant ribosomal proteins <203> (<203>, two plant ribosomal proteins of 14000 Da and 16000 Da can be phosphorylated by the Atpk1 protein kinase [379]) (Reversibility: ? <203> [379]) [379]
- P** ADP + phosphorylated plant ribosomal proteins
- S** ATP + protein <125> (<125>, autophosphorylation [245]; <125>, blue light-dependent autophosphorylating [244]) (Reversibility: ? <125> [244, 245]) [244, 245]
- P** ADP + phosphoprotein
- S** ATP + protein <205> (<205>, major autophosphorylation sites are Ser484 and Thr485 [382]) (Reversibility: ? <205> [382]) [382]
- P** ADP + phosphoprotein
- S** ATP + protein <238, 254, 258, 259, 268, 269, 273> (<238, 254, 258, 259, 268, 269, 273>, autophosphorylation [433, 462, 469, 481, 482, 487]; <269>, Ser916 is an in vivo autophosphorylation site [482]) (Reversibility: ? <238, 254, 258, 259, 268, 269, 273> [433, 462, 469, 481, 482, 487]) [433, 462, 469, 481, 482, 487]
- P** ADP + phosphoprotein
- S** ATP + protein <7, 44, 48, 53, 97> (<7, 53, 97>, autophosphorylation [8,110,178]; <44>, specifically phosphorylates the agonist-occupied forms of the β_2 -adrenergic receptor and related G protein-coupled receptors [101]; <44>, the enzyme mediates agonist-dependent phosphorylation of the β_2 -adrenergic and related G protein-coupled receptors [98]; <48>, specifically phosphorylates the agonist-occupied form of the β -adrenergic and related G protein-coupled receptors [106]) (Reversibility: ? <7, 44, 48, 53, 97> [8, 98, 101, 106, 110, 178]) [8, 98, 101, 106, 110, 178]
- P** ADP + phosphoprotein
- S** ATP + protein kinase C ζ <98> (<98>, phosphorylation and activation of protein kinase C ζ [179]) (Reversibility: ? <98> [179]) [179]
- P** ADP + phosphorylated protein kinase C ζ
- S** ATP + rhodopsin <181, 205, 206> (<181>, reaction only with GRK4 α isoform, no reaction with GRK4 β , GRK4 γ , GRK4 δ [344]; <205>, GRK5 phosphorylates rhodopsin in a light-dependent manner, phosphorylates the C-terminal tail region [382]) (Reversibility: ? <181,205,206> [344,382,384]) [344, 382, 384]
- P** ADP + phosphorylated rhodopsin
- S** ATP + rhodopsin <53, 76, 86, 96> (<53>, light-dependent deactivation of rhodopsin involves receptor phosphorylation that is mediated by the

highly specific protein kinases rhodopsin kinase [109]) (Reversibility: ? <53, 76, 86, 96> [109, 146, 147, 148, 177]) [109, 146, 147, 148, 177]

P ADP + phosphorylated rhodopsin

S ATP + ribosomal protein S6-(229-239) peptide analogue <272> (Reversibility: ? <272> [486]) [486]

P ADP + phosphorylated ribosomal protein S6-(229-239) peptide analogue

S Additional information <170, 204, 206, 210> (<170>, phorbol ester receptor/protein kinase [326]; <204>, preferably phosphorylates the *Saccharomyces cerevisiae* Pkc1p pseudosubstrate peptide and myelin basic protein, but not histones, protamine or dephosphorylated casein [381]; <206>, GRK6, but not other GRKs tested, incorporated tritium after incubation with [³H]palmitate in Sf9 and in COS-7 cells overexpressing the kinase [383]; <210>, interaction between RAC-PK and protein kinase C [390]) [326, 381, 383, 390]

P ?

Inhibitors

1-(5-isoquinolinesulfonyl)-2-methylpiperazine <110> [191]

1-oleoyl-2-acetyl glycerol <269> (<269>, inhibits phorbol ester binding [483]) [483]

Ro 31-8220 <114> [227]

calmodulin <181> (<181>, reaction of isoenzyme GRK4 α with rhodopsin, IC₅₀: 80 nM [344]) [344]

staurosporine <8, 114, 193, 194> [14, 227, 364]

Additional information <110> (<110>, insensitive to caspase-3 [191]) [191]

Additional information <243> (<243>, insensitive to PKC inhibitors known to interfere either with the regulatory or the catalytic domain [268]) [268]

Additional information <8> (<8>, modification and concomitant inactivation of the catalytic subunit of bovine heart cAMP-dependent protein kinase with affinity analogs of peptide substrates potentially capable of undergoing disulfide interchange with enzyme-bound sulfhydryl groups [12]) [12]

Cofactors/prosthetic groups

3-phosphoinositide <4, 98> (<4,98>, enzyme is dependent on [4,179]) [4, 179]

8-bromo-cAMP <66> (<66>, 50% activation at 0.00062 mM [130]) [130]

8-bromo-cGMP <66> (<66>, 50% activation at 0.00004 mM [130]) [130]

FMN <125> (<125>, apoprotein noncovalently binds FMN to form the holo-protein nph1 [244]) [244]

cAMP <1, 8, 9, 11, 12, 14, 15, 27, 34, 37, 40, 41, 52, 58, 61, 62, 66> (<1, 8, 9, 11, 12, 27, 37, 40, 41, 58, 61, 62>, dependent on [1, 3, 12, 13, 14, 17, 24, 25, 27, 28, 29, 30, 31, 32, 67, 68, 69, 89, 93, 116, 117, 118, 122, 123, 124]; <66>, 50% activation at 0.0117 mM, activation is not cooperative [130]) [1, 3, 12, 13, 14, 17, 24, 25, 27-32, 67-69, 84, 89, 93, 108, 116-118, 122-124, 130]

cGMP <6, 7, 30, 36, 57, 66, 70, 73, 84, 88, 97> (<6, 7, 30, 36, 57, 66, 70, 73, 84, 88, 97>, dependent on [6, 7, 8, 9, 10, 11, 67, 77, 78, 88, 115, 133, 134, 135, 140,

161, 166, 178]; <66>, cGMP, 0.010 mM, stimulated histone H2B phosphorylation by the DG1 protein kinase 20-fold [130]; <66>, 50% activation at 0.00019 mM, cooperative activation [130]) [6, 7, 8, 9, 10, 11, 67, 77, 78, 88, 115, 130, 133, 134, 135, 140, 161, 166, 178]

cIMP <66> (<66>, 50% activation at 0.0053 mM [130]) [130]

flavin <125> (<125>, LOV1 and LOV2 may be flavin-binding domains that regulate kinase activity in response to blue light-induced redox changes [246]; <125>, flavoprotein [246]) [245, 246]

Activating compounds

AMP <166> (<166>, dependent on AMP [315]) [315]

arachidonic acid <137> (<137>, alone or a combination of γ -linolenic acid and phosphatidylserine slightly enhances PKC ζ activity [268]) [268]

arachidonic acid <243> (<243>, slightly enhances PKC ζ activity [268]) [268]

cardiolipin <272> (<272>, activates [486]) [486]

diacylglycerol <137, 140, 145> (<140> activates [272]; <137>, plus phosphatidylserine, activates [268]; <145>, activity is dependent on [290]) [268, 272, 290]

diacylglycerol <146> (<146>, activity is dependent on [290]; <147>, independent of the presence of Ca^{2+} or diacylglycerol, when assayed with calf thymus H1 histone as a phosphate acceptor protein [292]) [290]

diacylglycerol <244> (<244>, dependent upon phosphatidylserine or diacylglycerol for maximal activation [441]) [441]

γ -linolenic acid <243> (<243>, a combination of γ -linolenic acid and phosphatidylserine slightly enhances PKC zeta activity [268]) [268]

gastrin <116> (<116>, physiological activator of PKD2 in human AGS-B cells stably transfected with the CCK(B)/gastrin receptor [190]) [190]

insulin <113> (<113>, activated endogenous protein kinase B α_1 2-fold in L6 myotubes, while after transfection into 293 cells PKB α is activated 20- and 50-fold in response to insulin and IGF-1 respectively. In both cells, the activation of PKB α is accompanied by its phosphorylation at Thr308 and Ser473 [221]) [221]

neurabin <165> (<165>, by way of its PDZ domain, the neuronal-specific neurabin may target p70(S6k) to nerve terminals [311]) [311]

pervanadate <113> (<113>, activation of Akt is associated with tyrosine phosphorylation of Akt [187]) [187]

phorbol dibutyrate <140> (<140>, activates [272]) [272]

phorbol esters <174> (<174>, bind to and stimulate the kinase activity of PKC-L [336]) [336]

phorbol esters <234> (<234>, stimulate [429]) [429]

phosphatidylserine <137> (<137>, plus diacylglycerol, activates [268]) [268]

phosphatidylserine <243> (<243>, a combination of γ -linolenic acid and phosphatidylserine slightly enhances PKC zeta activity [268]) [268]

phosphatidylserine <244, 260, 261> (<244>, dependent upon phosphatidylserine or diacylglycerol for maximal activation [441]; <260>, stimulates [471]; <260,261>, PKC Apl I requires much less phosphatidylserine for activation than does purified PKC Apl II [471,474]) [441, 471, 474]

phospholipid <110, 145> (<110>, strict dependence on the presence of phospholipids, [212]; <145>, activity is dependent on [290]) [212, 290]

phospholipid <146, 147> (<146>, activity is dependent on [290]; <147>, significantly dependent on phospholipid when assayed with calf thymus H1 histone as a phosphate acceptor protein [292]) [290]

phospholipid <275, 278> (<275>, dependent on [489]; <278>, stimulates [429]) [429, 489]

serum <113> (<113>, activation of Akt is associated with tyrosine phosphorylation of Akt [187]) [187]

Additional information <111, 113> (<111>, activation of serum-regulated and glucocorticoid-regulated protein kinase by agonists that activate phosphatidylinositol 3-kinase is mediated by 3-phosphoinositide-dependent protein kinase-1 and PDK2 [215]; <111>, activation of SGK by IGF-1 or hydrogen peroxide is initiated by a PtdIns(3,4,5)P₃-dependent activation of PDK2, which phosphorylates Ser422. This is followed by the PtdIns(3,4,5)P₃-independent phosphorylation at Thr256 that activates SGK, and is catalysed by PDK1 [216]; <113>, PKB α becomes phosphorylated and activated in insulin/IGF-1-stimulated cells by an upstream kinase(s) [221]) [215, 216, 221]

Additional information <243, 245> (<243>, cannot be activated by phorbol ester treatment of NIH 3T3 cells or insect cells, overexpressing the respective PKC isoenzyme [268]; <245>, transcriptionally regulated by serum and glucocorticoids in mammary epithelial cells [442]) [268, 442]

Additional information <7> (<7>, aminoterminal dimerization site of cGMP-dependent protein kinase and the autophosphorylation site, present in this part, control not only the activation of the enzyme but also the cooperative binding characteristics of the intact enzyme [8]) [8]

Metals, ions

Ca²⁺ <109> (<109>, Ca²⁺-mediated interactions between the two domains could contribute to enzyme activation as well as to the creation of a positively charged phosphatidylserine-binding site [198]; <110>, no activation [212]; <137>, activity is independent of Ca²⁺ [268]) [198]

Ca²⁺ <260> (<260>, enzyme is dependent on Ca²⁺ [471]; <260>, activates [472]) [471, 472]

Mg²⁺ <66> (<66>, maximal activation at 40-50 mM [130]) [130]

Additional information <202> (<202>, presence of only one cysteine-rich, zinc finger-like domain, absence of an apparent Ca(2+)-binding region [377]) [377]

K_m-Value (mM)

0.0036 <272> (S6-(229-239) peptide, <272>, enzyme activated by cardiolipin [486]) [486]

4 Enzyme Structure

Molecular weight

Additional information <8> (<8>, amino acid sequence of the catalytic subunit [15]) [15]

Subunits

? <116> (<116>, x * 105000, SDS-PAGE [190]) [190]

? <147, 165, 202, 203> (<203> x * 52554, calculation from nucleotide sequence [380]; <165>, x * 56160 [312]; <165>, x * 59109 [313]; <147>, x * 64000, SDS-PAGE [292]; <202>, x * 65000 [377]) [380, 313, 312, 292, 377]

? <238, 245, 248, 258, 272, 278> (<245>, x * 49000 [445]; <248>, x * 70000 [450]; <238>, x * 79000, SDS-PAGE [433]; <258>, x * 103925, calculation from nucleotide sequence [470]; <272>, x * 116000 [486]; <258>, x * 120000, SDS-PAGE [470]; <278>, x * 126000 [429]) [429, 433, 445, 450, 470, 486]

? <7, 30> (<7>, x * 76331 [10]; <30>, x * 77803, calculation from nucleotide sequence [78]) [10, 78]

Additional information <171> (<171> p70 S6 kinase α I with a calculated MW of 58946 Da consists of 525 amino acids, of which the last 502 residues are identical in sequence to the entire polypeptides encoded by the p70 S6 kinase α II with a calculated MW of 56153 Da. Both p70 S6 kinase polypeptides predicted by these cDNAs are present in p70 S6 kinase purified from rat liver, and each is thus expressed in vivo. The slightly longer α I polypeptide exhibits anomalously slow mobility on SDS-PAGE, migrating at an apparent MW of 90000 Da probably because of the presence of six consecutive Arg residues immediately following the initiator methionine [327]) [327]

Additional information <8, 14, 37, 58, 61> (<8>, MW of the catalytic subunit determined by amino acid sequence is 40580 Da [16]; <14>, three different genes encode the catalytic subunits of the cAMP-dependent protein kinase [30]; <37>, MW of the catalytic subunit C is 39000-41000 Da [89]; <58>, PkaC is a catalytic subunit of the Dictyostelium discoideum cAPK [116]; <61>, the product of pka1 is a catalytic subunit of protein kinase A [122]) [16, 30, 89, 116, 122]

Posttranslational modification

lipoprotein <206> (<206>, palmitoylation of GRK6 appears essential for membrane association [383]) [383]

phosphoprotein <109, 110, 113, 114, 116> (<113>, in SKOV3 ovarian carcinoma cells that exhibit high basal levels of Akt activity, Akt is tyrosine-phosphorylated in the basal state, and this phosphorylation is further enhanced by both pervanadate and insulin-like growth factor-1, Tyr474 is the site of tyrosine phosphorylation [187]; <116>, phorbol 12,13-dibutyrate in the presence of dioleoylphosphatidylserine stimulates the autophosphorylation of PKD2 in a synergistic fashion. Phorbol esters also stimulate autophosphorylation of PKD2 in intact cells, C-terminal Ser876 is an in vivo phosphorylation site within PKD2 that is correlated with the activation status of the kinase [190];

<109>, protein kinase C is processed by three phosphorylations. Firstly, trans-phosphorylation on the activation loop T500 renders it catalytically competent to autophosphorylate. Secondly, a subsequent autophosphorylation on the carboxyl terminus T641 maintains catalytic competence. Thirdly, a second autophosphorylation on the carboxyl terminus S660 regulates the enzyme's subcellular localization [199]; <109>, phosphorylation of Thr642 is an early event in the processing of newly synthesized protein kinase C β 1 and is essential for its activation [200]; <109>, processing by protein kinase C cannot occur until the enzyme is first phosphorylated by a protein kinase C kinase [201]; <109>, COOH-terminal autophosphorylation sites are critical for enzyme function and possibly subcellular localization in COS cells [202]; <110>, activation loop phosphorylation of PKC δ in response to serum stimulation of cells is PI 3-kinase-dependent and is enhanced by PDK1 coexpression [209]; <114>, two splice variants of protein kinase B γ have different regulatory capacity depending on the presence or absence of the regulatory phosphorylation site Ser472 in the carboxyl-terminal hydrophobic domain [225]; <114>, Akt3 is phosphorylated in response to insulin [226]; <114>, Akt-3 also possess a C-terminal 'tail' that contains a phosphorylation site Ser472 thought to be involved in the activation of Akt kinases. In addition to phosphorylation of Ser472, phosphorylation of Thr305 also appears to contribute to the activation of Akt-3 [227]; <114>, regulatory phosphorylation sites in the activation loop and in the C-terminal hydrophobic domain, Thr305 and Ser472, phosphorylation of both sites is required for full activity [228]) [187, 190, 199, 200, 201, 202, 209, 225, 226, 227, 228]

phosphoprotein <114, 286> (<114,286>, two splice variants of protein kinase B γ have different regulatory capacity depending on the presence or absence of the regulatory phosphorylation site Ser472 in the carboxyl-terminal hydrophobic domain, activation of PKB γ 1 requires phosphorylation at a single regulatory site Thr305 [225]; <114,286>, regulatory phosphorylation sites in the activation loop and in the C-terminal hydrophobic domain, Thr305 and Ser472 [225]) [225]

phosphoprotein <205> (<205>, major autophosphorylation sites are Ser484 and Thr485 [382]) [382]

phosphoprotein <7, 8> (<7>, autophosphorylation [8,9]; <8>, phosphate groups at Thr196 and Ser337 [16]) [8, 9, 16]

Additional information <109> (<109>, two types of complementary DNA clones for rat brain protein kinase C, these clones encode 671 and 673 amino acid sequences, which differ from each other only in the carboxyl-terminal regions of approximately 50 amino acid residues. This difference seems to result from alternative splicing [207]) [207]

Additional information <34> (<34>, unmyristylated C α ₂ may be essential for fertility in the male [85]) [85]

5 Isolation/Preparation/Mutation/Application

Source/tissue

3T3 cell <164> [310]

AGS <116> (<116>, AGS-B cell [190]) [190]

HL-60 cell <116> [190]

SKOV-3 cell <113> [187]

aorta <233> [428]

blood platelet <242> [437]

brain <109, 110, 111, 113, 135, 139, 140, 145> (<139>, in adult brain, the relative activities of α -, β I-, β II-, and γ -subspecies are roughly 16%, 8%, 55%, and 21% [271]) [191, 206, 207, 208, 211, 216, 224, 264, 270, 271, 290]

brain <146, 147, 148, 152, 181, 202, 205, 206> (<205>, very weak activity [382]) [281, 290, 292, 295, 345, 377, 382, 384]

brain <233, 241, 242, 245, 248, 268, 273, 285> (<245>, the cells which strongly express the *sgk* gene are in the deep layers of the cortex and in the corpus callosum. It is likely that the *sgk* transcript is expressed by oligodendrocytes after brain injury. Neurons in layers I and II of the cortex, lateroposterior and laterodorsal thalamic nucleus, and ventral posterolateral and posteromedial thalamic nucleus strongly express *sgk* mRNA at postnatal day 1 and day 7, but these neurons show no expression in fetal or adult brain. Induction of *sgk* gene may be associated with a series of axonal regenerations after brain injury, and in addition, the *sgk* gene may also play important roles in the development of particular groups of neurons in the postnatal brain [444]; <248>, different expression of the myotonin protein kinase gene in discrete areas of brain [449]) [428, 436, 437, 444, 449, 451, 481, 487, 500]

brain <38, 47, 49, 88> (<47,49>, synapses [104]; <47>, β -ARK mRNA is expressed intensely in the cerebellar granule cell layer and moderately in the hippocampal pyramidal cells and dentate granule cells. The neocortex and piriform cortex express it moderately to weakly, whereas the thalamus and hypothalamus express it weakly to faintly. No significant expression of the mRNA is detected in the caudate-putamen. Weak expression of β -ARK mRNA in several nuclei of the brainstem and in the spinal gray matter [105]) [90, 104, 105, 166]

cardiac muscle <8> [16]

cell culture <107, 111, 113> (<107>, NMuMG epithelial cell line [196]; <111>, hepatoma cell line [218]; <113>, cell lines MCF-7 and WI38 [223]; <113>, SKOV3 ovarian carcinoma cell [187]) [187, 196, 218, 223]

cell culture <160, 174, 180, 181, 202> (<160>, PR-17 cells and wild-type HL-60 cells [304]; <164>, Swiss 3T3 fibroblasts [310]; <174>, epidermoid carcinoma line A431 [336]; <164>, fibrosarcoma cell line [309]; <180>, ovarian carcinoma cell lines [342]; <180>, cell lines MCF-7 and WI38 [343]; <181>, spermatogonia cell line GC-1 spg [344]; <202>, insulin-secreting cell line RINm5F [377]) [304, 309, 310, 336, 342, 343, 344, 377]

cell culture <216, 241, 245, 268> (<216>, HeLa cells [400]; <241>, several hemopoietic tumor lines [436]; <245>, Con8.hd6 rat mammary tumor cell

line [445]; <268>, undifferentiated mouse embryonal carcinoma cell line P19, NIH 3T3 cells [481]) [400, 436, 445, 481]
 cell culture <13, 45, 52> (<45>, ovary cell line [26]; <52>, myoblast L6 cell line [108]) [26, 108]
 collecting duct <107> [195]
 colon <107> [194]
 colon <270> [484]
 embryo <37, 98> [89, 179]
 embryo <245> (<245>, embryo kidney 293 cells [442]) [442]
 erythroleukemia cell <283> [498]
 fibroblast <109, 113> [204, 224]
 fibroblast <70> (<70>, in normal diploid fibroblasts, the gene is constitutively expressed during cell-cycle and population doubling levels [133]) [133]
 fibrosarcoma cell line <164> [309]
 frontal cortex <243> [439]
 germ <181> [344]
 heart <113> [224]
 heart <233> [428]
 heart <174, 205, 206> (<205>, high activity [382]) [335, 336, 382, 384]
 heart <6, 8, 89> (<6>, expressed at a much higher level in newborn than in adult [6]; <89>, enhanced expression of β -adrenergic receptor kinase 1 in the hearts of cardiomyopathic Syrian hamsters, BIO53.58 [167]) [6, 12, 167]
 hematopoietic cell <110, 116> (<116>, hematopoietic stem cell and hematopoietic progenitor cells [229]) [212, 229]
 hematopoietic cell <242> (<242>, enzyme is predominantly expressed in [438]) [438]
 hepatoma cell <165> [314]
 hippocampus <258> [470]
 intestine <88> (<88>, intestinal mucosa [166]) [166]
 kidney <88> [166]
 kidney <233, 254, 268, 270, 285> [428, 462, 481, 484, 500]
 kidney <107, 110, 111, 122> (<122>, cortex [241]; <122>, low expression in vascular elements and high expression in tubule epithelium, highly expressed in proximal tubule, thick limb, and collecting duct [241]) [191, 194, 216, 241]
 kidney <205> (<205>, very weak activity [382]) [382]
 larva <37> [89]
 larva <190> (<190>, abundant at the earliest larval stage, but their relative concentrations decrease coordinately in late larvae [361]) [361]
 leukocyte <141> [277]
 leukocyte <44, 59> (<44,59>, peripheral blood leukocytes [99,120]) [99, 120]
 liver <233, 272> [428, 486]
 liver <111, 114> (<114>, not highly expressed [227]) [216, 227]
 liver <165, 171, 205, 206> (<205>, very weak activity [382]) [312, 327, 382, 384]
 lung <113> [224]
 lung <7> [9, 10]

lung <170, 174, 202, 205, 206> (<205>, high activity [382]) [326, 335, 336, 377, 382, 384]
lung <233, 241, 242, 245, 270> (<245>, high activity [445]) [428, 436, 437, 445, 484]
mammary epithelium <245> [196]
mature ovarian follicle <245> [443]
megakaryoblast <242> [437]
myeloid cell <110> (<110>, ABPL-3 myeloid tumor [212]) [212]
nervous system <260, 261> [472]
ovary <149, 150> [293]
ovary <110> (<110>, highly expressed [191]) [191]
ovary <245, 268> (<245>, high activity [445]) [445, 481]
pancreas <111> [216]
pancreas <206> [384]
pancreatic islet <202> [377]
pineal gland <76> [148]
placenta <206> [384]
placenta <216> [399]
placenta <30> [78]
retina <115> [189]
retina <44> [98]
retina <205> (<205>, high activity [382]) [382]
retinal cone <115> [189]
sensory cell <260, 261> [473]
skeletal muscle <206> [384]
skeletal muscle <114> (<114>, not highly expressed [227]) [227]
skeletal muscle <233, 242, 255> (<233>, predominantly expressed in [428]) [428, 437, 463]
skin <170, 174> [326, 335, 336]
small intestine <70> [134]
small intestine <233> [428]
sperm <34> [84, 85]
sperm <181> (<181>, GRK4 γ is the only detectable isoform in human sperm [344]) [344]
spermatozoon <166, 181> (<166>, transcripts encoding CAPL-B, an apparent member of the cyclic-nucleotide-regulated kinase subfamily in *Aplysia californica*, are found exclusively in the ovotestis and are concentrated in meiotic and postmeiotic spermatogenic cells. The CAPL-B polypeptide is present in mature spermatozoa [315]) [315, 344]
stomach <233> [428]
testis <98> [179]
testis <241, 268> [436, 481]
testis <40> [94]
testis <110> (<110>, highly expressed [191]) [191]
testis <181, 205, 210, 211> (<205>, very weak activity [382]) [344, 382, 390]
thymocyte <110> [191]
thymus <233, 245, 290> (<245>, high activity [445]) [428, 445, 504]

thyroid gland <270> [484]

tongue epithelium <205> [382]

trachea <7> (<7>, trachea smooth muscle [11]) [11]

uterus <233> [428]

Additional information <111, 114> (<111>, SGK3 is expressed in all tissues examined, but SGK2 mRNA is only present at significant levels in liver, kidney and pancreas and, at lower levels, in the brain [216]; <114>, expressed widely [227]) [216, 227]

Additional information <12, 37> (<12>, C α mRNA is widespread and highly expressed in brain, heart, adrenal gland, testis, lung, kidney, spleen and liver, whereas the C β mRNA is unevenly expressed in the brain and adrenal gland and in much lesser amounts in other tissues [27]; <37>, expressed at a low level in cytosolic and particulate compartments during embryogenesis. As the nematodes progress from late embryonic stages to the newly hatched, first larval stage, C subunit content increases 15fold. High levels of C subunits are observed in several subsequent larval and adult stages of development [89]) [27, 89]

Additional information <205> (<205>, most abundantly in lung, heart, retina, and lingual epithelium, but expressed very little in brain, liver, kidney, or testis [382]) [382]

Localization

cytoplasm <145> [286]

cytoplasm <272> [486]

cytoskeleton <190> (<190>, associated with [361]) [361]

cytosol <37> [89]

cytosol <145> (<145>, stimulation by phorbol ester causes weak translocation of δ III-GFP from the cytosol to the plasma membrane [286]) [286]

membrane <88> [166]

membrane <164, 190, 205> (<190>, associated with [361]; <205>, GRK5 protein does not undergo agonist-dependent translocation from cytosol to membranes as do β -adrenergic receptor kinase and rhodopsin kinase, but rather appears to associate with membranes constitutively [382]) [309, 361, 382]

membrane <114, 269, 286> (<269>, the enzyme contains membrane localization signals [483]; <114,286>, phosphorylation of the hydrophobic motif at the extreme C terminus of PKB γ may facilitate translocation of the kinase to the membrane [225]) [483, 225]

perinuclear space <164> [308]

plasma membrane <145> (<145>, stimulation by phorbol ester causes weak translocation of δ III-GFP from the cytosol to the plasma membrane [286]) [286]

ribosome <203> [379, 380]

ribosome <14> (<14>, 54 S subunit of the yeast mitochondrial ribosome [29]) [29]

Additional information <245> (<245>, treatment of rat mammary tumor cells with serum caused hyperphosphorylation of endogenous SGK, and promoted translocation to the nucleus [442]) [442]

Purification

- <48> (expression in COS 7 cells [106]) [106]
- <110> [212]
- <113> [224]
- <135> [264]
- <137> (partial purification of the PKC-zeta isoenzyme [268]) [268]
- <145> (partial [290]) [290]
- <146> (partial [290]) [290]
- <205> (recombinant enzyme [205]) [382]
- <243> (partial [268]) [268]
- <272> [486]
- <278> (partial [429]) [429]

Crystallization

- <8> (crystal structures of catalytic subunit of cAMP-dependent protein kinase in complex with isoquinolinesulfonyl protein kinase inhibitors 1-(5-isoquinolinesulfonyl)-2-methylpiperazine, N-[2-(methylamino)ethyl]-5-isoquinolinesulfonamide and N-[2-(*p*-bromocinnamylamino)ethyl]-5-isoquinolinesulfonamide [13]; crystal structure of the protein kinase catalytic subunit with staurosporine bound to the adenosine pocket [14]) [13, 14]
- <110> (crystal structure of the cys2 activator-binding domain of protein kinase C δ in complex with phorbol ester [210]) [210]
- <145> (crystal structure of PKC- δ C₂ domain. Structural elements unique to this C₂ domain include a helix and a protruding β hairpin which may contribute basic sequences to a membrane-interaction site [288]) [288]

Cloning

- <6> (expression in COS cells [6]) [6]
- <7> (cDNA of the two isoforms of bovine cGMP-dependent protein kinase [11]) [11]
- <9> (isolation of a full-length cDNA clone coding for C β 2 isoform of the bovine C-subunit [17]) [17]
- <11> [24]
- <12> (characterization of genomic clones coding for the C α and C β subunits [25]; isolation of a full-length cDNA clone encoding the C β catalytic subunit of cAMP-dependent protein kinase from a brain cDNA library [27]) [25, 27]
- <13> [26]
- <27> [68]
- <30> (characterization of the human gene encoding the type I α and type I β cGMP-dependent protein kinase [77]) [77]
- <33> [69]
- <34> [85]
- <36> [11]
- <37> [89]
- <38> (expression in COS-7 cells [90]) [90]
- <40> [93, 94]
- <41> [93]

- <44> (β ARK locus segregated with the long arm of chromosome 11, centromeric to 11q13 [98]; cloning and sequencing [99]) [98, 99]
- <52> (isolation of a full length cDNA clone encoding the C α type catalytic subunit of cAMP-dependent protein kinase [108]) [108]
- <53> [109]
- <58> (isolation of cDNA [117]; expression of the catalytic subunit DdPK3 in *Escherichia coli* [118]) [117, 118]
- <59> (expression in COS7 cells [120]) [120]
- <61> [122, 123]
- <62> [124]
- <66> (expressed in Sf9 cells [130]) [130]
- <70> [133, 134]
- <73> [77, 140]
- <76> [148]
- <84> (expression in COS-1 cells [161]) [161]
- <86> [148]
- <88> [166]
- <89> [167]
- <95> [176]
- <98> [179]
- <109> (expression of mutant enzymes in COS cells [201]; wild-type and mutant enzymes overexpressed in COS cells [202]; isolation of cloned mouse protein kinase C β -II cDNA [203]; expression in COS cells [208]) [201, 202, 206, 208]
- <109> (isolation of cDNA clones encoding protein kinase C [205]) [205]
- <110> (PKC δ II expressed in COS-1 cells [191]) [191]
- <111> [218]
- <113> (characterization of the gene [192]) [192]
- <113> (expression in COS cells [224]) [222, 223, 224]
- <114> [227, 228]
- <115> (GRK7 [189]) [189]
- <116> [190, 229]
- <122> [241]
- <124> [242]
- <137> (expression in recombinant baculovirus-infected insect cells [268]) [268]
- <138> (structure and nucleotide sequence of a *Drosophila melanogaster* protein kinase C gene [269]) [269]
- <140> (cDNA sequence encoding mouse PKC- γ isolated from a C57BL/6 brain cDNA library [273]) [273]
- <140> (expression in COS cells [208]) [208]
- <141> (the 5' segment of the gene for protein kinase C β is cloned from a human leukocyte genomic library in EMBL3 bacteriophage [277]) [276, 277]
- <145> (cDNA cloning of an alternative splicing variant of protein kinase C δ , expression of truncated form of PKC δ III fused to green fluorescent protein in CHO-K1 cells [286]; expressed in COS 7 cells [290]) [286, 290, 291]
- <203> (isolation of cDNA [378]) [378]

- <216> [399, 400]
- <234> [429]
- <236> [222, 224]
- <238> (expression in COS1 cells transfected with an nPKC theta cDNA expression plasmid [433]) [433]
- <239> (protein kinase C- ϵ E1 and E2, expression in Sf9 cells, the recombinant protein displays protein kinase C activity and phorbol ester binding activity [434]) [434]
- <241> (expressed in insect cells via a baculovirus expression vector, a 75000 Da protein is synthesized which, unlike other PKC isoforms, does not bind phorbol ester, even at very high concentrations [436]) [436]
- <242> (expression in COS cells [437]) [437, 438]
- <243> (expression in recombinant baculovirus-infected insect cells, overexpression in NIH 3T3 cells or insect cells [268]; isolation of cDNA [439]) [268, 439]
- <244> (expression in the baculovirus insect-cell expression system [441]) [441]
- <248> (expression in COS-1 cells [450]) [450]
- <255> (expression in COS cells [463]) [400, 463]
- <256> [400]
- <258> (expression in COS1 cells [469]; expression in COS7 cells [470]) [469, 470]
- <259> (expression in COS1 cells [469]) [469]
- <260> [473]
- <261> [473]
- <267> (isolation of cDNA [480]) [480]
- <268> (COS cells transfected with the PKC lambda expression plasmid [481]) [481]
- <269> (bacterially expressed catalytic domain of PKD [483]) [483]
- <273> [487]
- <274> (DNA sequences encoding the rat Rsk-1 S6 kinase modified by insertion of a peptide epitope at the polypeptide aminotermminus, expressed in COS cells [488]) [488]
- <275> (expression in COS cells [489]) [489]
- <278> [429]
- <283> [498, 499]
- <286> [228]

Engineering

- M24L <171> (<171>, mutation in the α I polypeptide suppresses synthesis of the α II polypeptides [327]) [327]
- M24T <171> (<171>, mutation in the α I polypeptide suppresses synthesis of the α II polypeptides [327]) [327]
- S442X <245> (<245>, mutation of putative phosphorylation sites at Thr256 and Ser422 inhibited SGK activation [442]) [442]
- S79D <97> (<97>, replacement of an autophosphorylated Ser79 of cGKI β with an aspartic acid results in a mutant kinase with constitutive kinase ac-

tivity in vitro and in vivo. The cGKI β S79D mutant localized to the cytoplasm and is only a weak activator of CRE-dependent gene transcription [178]) [178]

T256A <107> (<107>, the conditional IPTG inducible expression of wild type Sgk, but not of the kinase dead mutant T256A Sgk, protects Con8 mammary epithelial tumor cells from serum starvation-induced apoptosis [181]) [181]

T256A/S422A <107> (<107>, unlike the wild-type enzyme the phosphorylation site mutant has no effect on cell survival [181]) [181]

T256X <245> (<245>, mutation of putative phosphorylation sites at Thr256 and Ser422 inhibited SGK activation [442]) [442]

T474F <113> (<113>, 55% inhibition of Akt activation [187]) [187]

T500E <109> (<109>, expression as a catalytically active protein kinase C in COS cells [201]) [201]

T500V <109> (<109>, expression as a catalytically inactive protein kinase C in COS cells [201]) [201]

Additional information <114> (<114>, phosphorylation of Ser472 and phosphorylation of Thr305 appears to contribute to the activation of Akt-3 because mutation of both these residues to aspartate increases the catalytic activity of Akt-3, whereas mutation to alanine inhibits activation [227]) [227]

Additional information <185> (<185>, mutant, gprk2(6936) disrupts expression of a putative member of the GRK family, the G protein-coupled receptor kinase 2 gene Gprk2. This mutation affects Gprk2 gene expression in the ovaries and renders mutant females sterile. The mutant eggs contain defects in several anterior eggshell structures that are produced by specific subsets of migratory follicle cells. In addition, rare eggs that become fertilized display severe defects in embryogenesis [349]) [349]

Additional information <248> (<248>, a CTG triplet repeat undergoes expansion in myotonic dystrophy patients. This sequence is highly variable in the normal population, unaffected individuals have been 5 and 27 copies, myotonic dystrophy patients are minimally affected have at least 50 repeats, more severely affected patients have expansion of the repeat containing segment up to several kilobase pairs [454]) [454]

References

- [1] Meyer, E.; Butler, A.; Dubrana, K.; Duharcourt, S.; Caron, F.: Sequence-specific epigenetic effects of the maternal somatic genome on developmental rearrangements of the zygotic genome in *Paramecium primaurelia*. *Mol. Cell. Biol.*, **17**, 3589-3599 (1997)
- [2] Verde, F.; Wiley, D.J.; Nurse, P.: Fission yeast orb6, a Ser/Thr protein kinase related to mammalian rho kinase and myotonic dystrophy kinase, is required for maintenance of cell polarity and coordinates cell morphogenesis with the cell cycle. *Proc. Natl. Acad. Sci. USA*, **95**, 7526-7531 (1998)
- [4] Stephens, L.; Anderson, K.; Stokoe, D.; Erdjument-Bromage, H.; Painter, G.F.; et al.: Protein kinase B kinases that mediate phosphatidylinositol

- 3,4,5-trisphosphate-dependent activation of protein kinase B. *Science*, **279**, 710-714 (1998)
- [6] Kumar, R.; Joyner, R.W.; Komalavilas, P.; Lincoln, T.M.: Analysis of expression of cGMP-dependent protein kinase in rabbit heart cells. *J. Pharmacol. Exp. Ther.*, **291**, 967-975 (1999)
- [7] Hashimoto, E.; Takio, K.; Krebs, E.G.: Amino acid sequence at the ATP-binding site of cGMP-dependent protein kinase. *J. Biol. Chem.*, **257**, 727-733 (1982)
- [8] Heil, W.G.; Landgraf, W.; Hofmann, F.: A catalytically active fragment of cGMP-dependent protein kinase. Occupation of its cGMP-binding sites does not affect its phosphotransferase activity. *Eur. J. Biochem.*, **168**, 117-121 (1987)
- [9] Takio, K.; Smith, S.B.; Walsh, K.A.; Krebs, E.G.; Titani, K.: Amino acid sequence around a "hinge" region and its "autophosphorylation" site in bovine Lung cGMP-dependent protein kinase. *J. Biol. Chem.*, **258**, 5531-5536 (1983)
- [10] Takio, K.; Wade, R.D.; Smith, S.B.; Krebs, E.G.; Walsh, K.A.; Titani, K.: Guanosine cyclic 3',5'-phosphate dependent protein kinase, a chimeric protein homologous with two separate protein families. *Biochemistry*, **23**, 4207-4218 (1984)
- [11] Wernet, W.; Flockerzi, V.; Hofmann, F.: The cDNA of the two isoforms of bovine cGMP-dependent protein kinase. *FEBS Lett.*, **251**, 191-196 (1989)
- [12] Bramson, H.N.; Thomas, N.; Matsueda, R.; Nelson, N.C.; Taylor, S.S.; Kaiser, E.T.: Modification of the catalytic subunit of bovine heart cAMP-dependent protein kinase with affinity labels related to peptide substrates. *J. Biol. Chem.*, **257**, 10575-10581 (1982)
- [13] Engh, R.A.; Girod, A.; Kinzel, V.; Huber, R.; Bossemeyer, D.: Crystal structures of catalytic subunit of cAMP-dependent protein kinase in complex with isoquinolinesulfonyl protein kinase inhibitors H7, H8, and H89. Structural implications for selectivity. *J. Biol. Chem.*, **271**, 26157-26164 (1996)
- [14] Prade, L.; Engh, R.A.; Girod, A.; Kinzel, V.; Huber, R.; Bossemeyer, D.: Staurosporine-induced conformational changes of cAMP-dependent protein kinase catalytic subunit explain inhibitory potential. *Structure*, **5**, 1627-1637 (1997)
- [15] Shoji, S.; Ericsson, L.H.; Walsh, K.A.; Fischer, E.H.; Titani, K.: Amino acid sequence of the catalytic subunit of bovine type II adenosine cyclic 3',5'-phosphate dependent protein kinase. *Biochemistry*, **22**, 3702-3709 (1983)
- [16] Shoji, S.; Parmelee, D.C.; Wade, R.D.; Kumar, S.; Ericsson, L.H.; Walsh, K.A.; Neurath, H.; Long, G.L.; Demaille, J.G.; Fischer, E.H.; Titani, K.: Complete amino acid sequence of the catalytic subunit of bovine cardiac muscle cyclic AMP-dependent protein kinase. *Proc. Natl. Acad. Sci. USA*, **78**, 848-851 (1981)
- [17] Wiemann, S.; Kinzel, V.; Pyerin, W.: Cloning of the C α catalytic subunit of the bovine cAMP-dependent protein kinase. *Biochim. Biophys. Acta*, **1171**, 93-96 (1992)

- [24] Showers, M.O.; Maurer, R.A.: A cloned bovine cDNA encodes an alternate form of the catalytic subunit of cAMP-dependent protein kinase. *J. Biol. Chem.*, **261**, 16288-16291 (1986)
- [25] Chrivia, J.C.; Uhler, M.D.; McKnight, G.S.: Characterization of genomic clones coding for the C α and C β subunits of mouse cAMP-dependent protein kinase. *J. Biol. Chem.*, **263**, 5739-5744 (1988)
- [26] Howard, P.; Day, K.H.; Kim, K.E.; Richardson, J.; Thomas, J.; Abraham, I.; Fleischmann, R.D.; Gottesman, M.M.; Maurer, R.A.: Decreased catalytic subunit mRNA levels and altered catalytic subunit mRNA structure in a cAMP-resistant Chinese hamster ovary cell line. *J. Biol. Chem.*, **266**, 10189-10195 (1991)
- [27] Shuntoh, H.; Sakamoto, N.; Matsuyama, S.; Saitoh, M.; Tanaka, C.: Molecular structure of the C β catalytic subunit of rat cAMP-dependent protein kinase and differential expression of C α and C β isoforms in rat tissues and cultured cells. *Biochim. Biophys. Acta*, **1131**, 175-180 (1992)
- [28] Uhler, M.D.; Chrivia, J.C.; McKnight, G.S.: Evidence for a second isoform of the catalytic subunit of cAMP-dependent protein kinase. *J. Biol. Chem.*, **261**, 15360-15363 (1986)
- [29] Fearon, K.; Mason, T.L.: Structure and function of MRP20 and MRP49, the nuclear genes for two proteins of the 54 S subunit of the yeast mitochondrial ribosome. *J. Biol. Chem.*, **267**, 5162-5170 (1992)
- [30] Toda, T.; Cameron, S.; Sass, P.; Zoller, M.; Wigler, M.: Three different genes in *S. cerevisiae* encode the catalytic subunits of the cAMP-dependent protein kinase. *Cell*, **50**, 277-287 (1987)
- [31] Vandenbol, M.; Bolle, P.A.; Dion, C.; Portetelle, D.; Hilger, F.: Sequencing and analysis of a 20.5 kb DNA segment located on the left arm of yeast chromosome XI. *Yeast*, **10**, S25-33 (1994)
- [32] Cannon, J.F.; Tatchell, K.: Characterization of *Saccharomyces cerevisiae* genes encoding subunits of cyclic AMP-dependent protein kinase. *Mol. Cell. Biol.*, **7**, 2653-2663 (1987)
- [33] Lisziewicz, J.; Godany, A.; Forster, H.H.; Kuntzel, H.: Isolation and nucleotide sequence of a *Saccharomyces cerevisiae* protein kinase gene suppressing the cell cycle start mutation *cdc25*. *J. Biol. Chem.*, **262**, 2549-2553 (1987)
- [34] Bussey, H.; Storms, R.K.; Ahmed, A.; Albermann, K.; et al.: The nucleotide sequence of *Saccharomyces cerevisiae* chromosome XVI. *Nature*, **387**, 103-105 (1997)
- [35] Ohno, S.; Aoshima, M.; Matsumoto, S.; Yahara, I.; Suzuki, K.: A yeast gene coding for a putative protein kinase homologous to *cdc25* suppressing protein kinase. *FEBS Lett.*, **222**, 279-285 (1987)
- [54] di Blasi, F.; Carra, E.; de Vendittis, E.; Masturzo, P.; Burderi, E.; Lambri-noudaki, I.; Mirisola, M.G.; Seidita, G.; Fasano, O.: The SCH9 protein kinase mRNA contains a long 5' leader with a small open reading frame. *Yeast*, **9**, 21-32 (1993)
- [55] Johnston, M.; Andrews, S.; Brinkman, R.; Cooper, J.; Ding, H.; Dover, J.; Du, Z.; Favello, A.; Fulton, L.; Gattung, S.; et al.: Complete nucleotide se-

- quence of *Saccharomyces cerevisiae* chromosome VIII. *Science*, **265**, 2077-2082 (1994)
- [56] Toda, T.; Cameron, S.; Sass, P.; Wigler, M.: SCH9, a gene of *Saccharomyces cerevisiae* that encodes a protein distinct from, but functionally and structurally related to, cAMP-dependent protein kinase catalytic subunits. *Genes Dev.*, **2**, 517-527 (1988)
- [67] Adams, M.D.; Celniker, S.E.; Holt, R.A.; Evans, C.A.; Gocayne, J.D.; et al.: The genome sequence of *Drosophila melanogaster*. *Science*, **287**, 2185-2195 (2000)
- [68] Foster, J.L.; Higgins, G.C.; Jackson, F.R.: Cloning, sequence, and expression of the *Drosophila* cAMP-dependent protein kinase catalytic subunit gene. *J. Biol. Chem.*, **263**, 1676-1681 (1988)
- [69] Kalderon, D.; Rubin, G.M.: Isolation and characterization of *Drosophila* cAMP-dependent protein kinase genes. *Genes Dev.*, **2**, 1539-1556 (1988)
- [77] Orstavik, S.; Natarajan, V.; Tasken, K.; Jahnsen, T.; Sandberg, M.: Characterization of the human gene encoding the type I α and type I β cGMP-dependent protein kinase (PRKG1). *Genomics*, **42**, 311-318 (1997)
- [78] Sandberg, M.; Natarajan, V.; Ronander, I.; Kalderon, D.; Walter, U.; Lohmann, S.M.; Jahnsen, T.: Molecular cloning and predicted full-length amino acid sequence of the type I β isozyme of cGMP-dependent protein kinase from human placenta. Tissue distribution and developmental changes in rat. *FEBS Lett.*, **255**, 321-329 (1989)
- [84] Agustin, J.T.; Wilkerson, C.G.; Witman, G.B.: The unique catalytic subunit of sperm cAMP-dependent protein kinase is the product of an alternative $C\alpha$ mRNA expressed specifically in spermatogenic cells. *Mol. Biol. Cell*, **11**, 3031-3044 (2000)
- [85] Desseyn, J.L.; Burton, K.A.; McKnight, G.S.: Expression of a nonmyristylated variant of the catalytic subunit of protein kinase A during male germ-cell development. *Proc. Natl. Acad. Sci. USA*, **97**, 6433-6438 (2000)
- [86] Maldonado, F.; Hanks, S.K.: A cDNA clone encoding human cAMP-dependent protein kinase catalytic subunit C α . *Nucleic Acids Res.*, **16**, 8189-8190 (1988)
- [88] Ruth, P.; Pfeifer, A.; Kamm, S.; Klatt, P.; Dostmann, W.R.; Hofmann, F.: Identification of the amino acid sequences responsible for high affinity activation of cGMP kinase I α . *J. Biol. Chem.*, **272**, 10522-10528 (1997)
- [89] Gross, R.E.; Bagchi, S.; Lu, X.; Rubin, C.S.: Cloning, characterization, and expression of the gene for the catalytic subunit of cAMP-dependent protein kinase in *Caenorhabditis elegans*. Identification of highly conserved and unique isoforms generated by alternative splicing. *J. Biol. Chem.*, **265**, 6896-6907 (1990)
- [90] Benovic, J.L.; DeBlasi, A.; Stone, W.C.; Caron, M.G.; Lefkowitz, R.J.: β -adrenergic receptor kinase: primary structure delineates a multigene family. *Science*, **246**, 235-240 (1989)
- [93] Beebe, S.J.; Oyen, O.; Sandberg, M.; Froysa, A.; Hansson, V.; Jahnsen, T.: Molecular cloning of a tissue-specific protein kinase (C γ) from human testis—representing a third isoform for the catalytic subunit of cAMP-dependent protein kinase. *Mol. Endocrinol.*, **4**, 465-475 (1990)

- [94] Reinton, N.; Haugen, T.B.; Orstavik, S.; Skalhogg, B.S.; Hansson, V.; Jahn-
sen, T.; Tasken, K.: The gene encoding the C γ catalytic subunit of cAMP-
dependent protein kinase is a transcribed retroposon. *Genomics*, **49**, 290-
297 (1998)
- [98] Benovic, J.L.; Stone, W.C.; Huebner, K.; Croce, C.; Caron, M.G.; Lefkowitz,
R.J.: cDNA cloning and chromosomal localization of the human β -adre-
nergic receptor kinase. *FEBS Lett.*, **283**, 122-126 (1991)
- [99] Chuang, T.T.; Sallese, M.; Ambrosini, G.; Parruti, G.; De Blasi, A.: High
expression of β -adrenergic receptor kinase in human peripheral blood
leukocytes. Isoproterenol and platelet activating factor can induce kinase
translocation. *J. Biol. Chem.*, **267**, 6886-6892 (1992)
- [100] Fushman, D.; Najmabadi-Haske, T.; Cahill, S.; Zheng, J.; LeVine, H.; Cow-
burn, D.: The solution structure and dynamics of the pleckstrin homology
domain of G protein-coupled receptor kinase 2 (β -adrenergic receptor ki-
nase 1). A binding partner of G $\beta\gamma$ subunits. *J. Biol. Chem.*, **273**, 2835-2843
(1998)
- [101] Penn, R.B.; Benovic, J.L.: Structure of the human gene encoding the β -
adrenergic receptor kinase. *J. Biol. Chem.*, **269**, 14924-14930 (1994)
- [104] Arriza, J.L.; Dawson, T.M.; Simerly, R.B.; Martin, L.J.; Caron, M.G.; Snyder,
S.H.; Lefkowitz, R.J.: The G-protein-coupled receptor kinases β ARK1 and
 β ARK2 are widely distributed at synapses in rat brain. *J. Neurosci.*, **12**,
4045-4055 (1992)
- [105] Owada, Y.; Watanabe, M.; Kondo, H.: Localization of mRNA for β -adre-
nergic receptor kinase in the brain of adult rats. *Neurosci. Lett.*, **144**, 9-
13 (1992)
- [106] Benovic, J.L.; Onorato, J.J.; Arriza, J.L.; Stone, W.C.; Lohse, M.; Jenkins,
N.A.; Gilbert, D.J.; Copeland, N.G.; Caron, M.G.; Lefkowitz, R.J.: Cloning,
expression, and chromosomal localization of β -adrenergic receptor kinase
2. A new member of the receptor kinase family. *J. Biol. Chem.*, **266**, 14939-
14946 (1991)
- [108] Wiemann, S.; Voss, H.; Kinzel, V.; Pyerin, W.: Rat C α catalytic subunit of
the cAMP-dependent protein kinase: cDNA sequence and evidence that it
is the only isoform expressed in myoblasts. *Biochim. Biophys. Acta*, **1089**,
254-256 (1991)
- [109] Lorenz, W.; Inglese, J.; Palczewski, K.; Onorato, J.J.; Caron, M.G.; Lefko-
witz, R.J.: The receptor kinase family: primary structure of rhodopsin ki-
nase reveals similarities to the β -adrenergic receptor kinase. *Proc. Natl.*
Acad. Sci. USA, **88**, 8715-8719 (1991)
- [110] Palczewski, K.; Buczylo, J.; Van Hooser, P.; Carr, S.A.; Huddleston, M.J.;
Crabb, J.W.: Identification of the autophosphorylation sites in rhodopsin
kinase. *J. Biol. Chem.*, **267**, 18991-18998 (1992)
- [115] Kalderon, D.; Rubin, G.M.: cGMP-dependent protein kinase genes in *Dro-*
sophila. *J. Biol. Chem.*, **264**, 10738-10748 (1989)
- [116] Anjard, C.; Etchebehere, L.; Pinaud, S.; Veron, M.; Reymond, C.D.: An
unusual catalytic subunit for the cAMP-dependent protein kinase of *Dic-*
tyostelium discoideum. *Biochemistry*, **32**, 9532-9538 (1993)

- [117] Burki, E.; Anjard, C.; Scholder, J.C.; Reymond, C.D.: Isolation of two genes encoding putative protein kinases regulated during *Dictyostelium discoideum* development. *Gene*, **102**, 57-65 (1991)
- [118] Mann, S.K.; Yonemoto, W.M.; Taylor, S.S.; Firtel, R.A.: DdPK3, which plays essential roles during *Dictyostelium* development, encodes the catalytic subunit of cAMP-dependent protein kinase. *Proc. Natl. Acad. Sci. USA*, **89**, 10701-10705 (1992)
- [119] Dunham, I.; Shimizu, N.; Roe, B.A.; Chisoe, S.; Hunt, A.R.; Collins, J.E.; Bruskiewich, R.; Beare, D.M.; Clamp, M.; Smink, L.J.; Ainscough, R.; Almeida, J.P.; Babbage, A.; Bagguley, C.; Bailey, J.; Barlow, K.; Bates, K.N.; Beasley, O.; Bird, C.P.; Blakey, S.; Bridgeman, A.M.; Buck, D.; Burgess, J.; Burrill, W.D.; O'Brien, K.P.; et al.: The DNA sequence of human chromosome 22. *Nature*, **402**, 489-495 (1999)
- [120] Parruti, G.; Ambrosini, G.; Sallese, M.; De Blasi, A.: Molecular cloning, functional expression and mRNA analysis of human β -adrenergic receptor kinase 2. *Biochem. Biophys. Res. Commun.*, **190**, 475-481 (1993)
- [122] Maeda, T.; Watanabe, Y.; Kunitomo, H.; Yamamoto, M.: Cloning of the *pka1* gene encoding the catalytic subunit of the cAMP-dependent protein kinase in *Schizosaccharomyces pombe*. *J. Biol. Chem.*, **269**, 9632-9637 (1994)
- [123] Yu, G.; Li, J.; Young, D.: The *Schizosaccharomyces pombe pka1* gene, encoding a homolog of cAMP-dependent protein kinase. *Gene*, **151**, 215-220 (1994)
- [124] Jung, S.; Hoffmann, R.; Rodriguez, P.H.; Mutzel, R.; Hofer, H.W.: The catalytic subunit of cAMP-dependent protein kinase from *Ascaris suum*. The cloning and structure of a novel subtype of protein kinase A. *Eur. J. Biochem.*, **232**, 111-117 (1995)
- [125] Jin, M.; Fujita, M.; Culley, B.M.; Apolinario, E.; Yamamoto, M.; Maundrell, K.; Hoffman, C.S.: *sck1*, a high copy number suppressor of defects in the cAMP-dependent protein kinase pathway in fission yeast, encodes a protein homologous to the *Saccharomyces cerevisiae* SCH9 kinase. *Genetics*, **140**, 457-467 (1995)
- [130] Foster, J.L.; Higgins, G.C.; Jackson, F.R.: Biochemical properties and cellular localization of the *Drosophila* DG1 cGMP-dependent protein kinase. *J. Biol. Chem.*, **271**, 23322-23328 (1996)
- [133] Fujii, M.; Ogata, T.; Takahashi, E.; Yamada, K.; Nakabayashi, K.; Oishi, M.; Ayusawa, D.: Expression of the human cGMP-dependent protein kinase II gene is lost upon introduction of SV40 T antigen or immortalization in human cells. *FEBS Lett.*, **375**, 263-267 (1995)
- [134] Orstavik, S.; Solberg, R.; Tasken, K.; Nordahl, M.; Altherr, M.R.; Hansson, V.; Jahnsen, T.; Sandberg, M.: Molecular cloning, cDNA structure, and chromosomal localization of the human type II cGMP-dependent protein kinase. *Biochem. Biophys. Res. Commun.*, **220**, 759-765 (1996)
- [135] Witczak, O.; Orstavik, S.; Natarajan, V.; Frengen, E.; Jahnsen, T.; Sandberg, M.: Characterization of the gene encoding the human type II cGMP-dependent protein kinase. *Biochem. Biophys. Res. Commun.*, **245**, 113-119 (1998)

- [140] Tamura, N.; Itoh, H.; Ogawa, Y.; Nakagawa, O.; Harada, M.; Chun, T.H.; Suga, S.; Yoshimasa, T.; Nakao, K.: cDNA cloning and gene expression of human type I α cGMP-dependent protein kinase. *Hypertension*, **27**, 552-557 (1996)
- [146] Khani, S.C.; Abitbol, M.; Yamamoto, S.; Maravic-Magovcevic, I.; Dryja, T.P.: Characterization and chromosomal localization of the gene for human rhodopsin kinase. *Genomics*, **35**, 571-576 (1996)
- [147] Yamamoto, S.; Sippel, K.C.; Berson, E.L.; Dryja, T.P.: Defects in the rhodopsin kinase gene in the Oguchi form of stationary night blindness. *Nat. Genet.*, **15**, 175-178 (1997)
- [148] Zhao, X.; Haeseleer, F.; Fariss, R.N.; Huang, J.; Baehr, W.; Milam, A.H.; Palczewski, K.: Molecular cloning and localization of rhodopsin kinase in the mammalian pineal. *Vis. Neurosci.*, **14**, 225-232 (1997)
- [161] Uhler, M.D.: Cloning and expression of a novel cyclic GMP-dependent protein kinase from mouse brain. *J. Biol. Chem.*, **268**, 13586-13591 (1993)
- [165] Tokumitsu, H.; Enslin, H.; Soderling, T.R.: Characterization of a Ca²⁺/calmodulin-dependent protein kinase cascade. Molecular cloning and expression of calcium/calmodulin-dependent protein kinase kinase. *J. Biol. Chem.*, **270**, 19320-19324 (1995)
- [166] Jarchau, T.; Hausler, C.; Markert, T.; Pohler, D.; Vanderkerckhove, J.; De Jonge, H.R.; Lohmann, S.M.; Walter, U.: Cloning, expression, and in situ localization of rat intestinal cGMP-dependent protein kinase II. *Proc. Natl. Acad. Sci. USA*, **91**, 9426-9430 (1994)
- [167] Urasawa, K.; Yoshida, I.; Takagi, C.; Onozuka, H.; Mikami, T.; Kawaguchi, H.; Kitabatake, A.: Enhanced expression of β -adrenergic receptor kinase 1 in the hearts of cardiomyopathic Syrian hamsters, BIO53.58. *Biochem. Biophys. Res. Commun.*, **219**, 26-30 (1996)
- [176] Alessi, D.R.; Deak, M.; Casamayor, A.; Caudwell, F.B.; Morrice, N.; Norman, D.G.; Gaffney, P.; Reese, C.B.; MacDougall, C.N.; Harbison, D.; Ashworth, A.; Bownes, M.: 3-Phosphoinositide-dependent protein kinase-1 (PDK1): structural and functional homology with the *Drosophila* DSTPK61 kinase. *Curr. Biol.*, **7**, 776-789 (1997)
- [177] Chen, C.K.; Burns, M.E.; Spencer, M.; Niemi, G.A.; Chen, J.; Hurley, J.B.; Baylor, D.A.; Simon, M.I.: Abnormal photoresponses and light-induced apoptosis in rods lacking rhodopsin kinase. *Proc. Natl. Acad. Sci. USA*, **96**, 3718-3722 (1999)
- [178] Collins, S.P.; Uhler, M.D.: Cyclic AMP- and cyclic GMP-dependent protein kinases differ in their regulation of cyclic AMP response element-dependent gene transcription. *J. Biol. Chem.*, **274**, 8391-8404 (1999)
- [179] Dong, L.Q.; Zhang, R.B.; Langlais, P.; He, H.; Clark, M.; Zhu, L.; Liu, F.: Primary structure, tissue distribution, and expression of mouse phosphoinositide-dependent protein kinase-1, a protein kinase that phosphorylates and activates protein kinase Czeta. *J. Biol. Chem.*, **274**, 8117-8122 (1999)
- [180] Boeckmann, B.; Bairoch, A.; Apweiler, R.; Blatter, M.C.; Estreicher, A.; Gasteiger, E.; Martin M.J.; Michoud, K.; O'Donovan, C.; Phan, I.; Pilbout,

- S.; Schneider, M.: The SWISS-PROT protein knowledgebase and its supplement TrEMBL. *Nucleic Acids Res.*, **31**, 365-370 (2003)
- [181] Leong, M.L.; Maiyar, A.C.; Kim, B.; O'Keefe, B.A.; Firestone, G.L.: Expression of the serum- and glucocorticoid-inducible protein kinase, Sgk, is a cell survival response to multiple types of environmental stress stimuli in mammary epithelial cells. *J. Biol. Chem.*, **278**, 5871-5882 (2003)
- [183] Saijo, K.; Mecklenbrauker, I.; Santana, A.; et al.: Protein kinase C β controls nuclear factor kappaB activation in B cells through selective regulation of the IkappaB kinase α . *J. Exp. Med.*, **195**, 1647-1652 (2002)
- [184] Mecklenbrauker, I.; Saijo, K.; Zheng, N.Y.; Leitges, M.; Tarakhovsky, A.: Protein kinase C δ controls self-antigen-induced B-cell tolerance. *Nature*, **416**, 860-865 (2002)
- [185] Gamper, N.; Fillon, S.; Feng, Y.; Friedrich, B.; Lang, P.A.; Henke, G.; Huber, S.M.; Kobayashi, T.; Cohen, P.; Lang, F.: K(+) channel activation by all three isoforms of serum- and glucocorticoid-dependent protein kinase SGK. *Pflugers Arch.*, **445**, 60-66 (2002)
- [187] Conus, N.M.; Hannan, K.M.; Cristiano, B.E.; Hemmings, B.A.; Pearson, R.B.: Direct identification of tyrosine 474 as a regulatory phosphorylation site for the Akt protein kinase. *J. Biol. Chem.*, **277**, 38021-38028 (2002)
- [188] Wiemann, S.; Weil, B.; Wellenreuther, R.; et al: Toward a catalog of human genes and proteins: sequencing and analysis of 500 novel complete protein coding human cDNAs. *Genome Res.*, **11**, 422-435 (2001)
- [189] Weiss, E.R.; Ducceschi, M.H.; Horner, T.J.; Li, A.; Craft, C.M.; Osawa, S.: Species-specific differences in expression of G-protein-coupled receptor kinase (GRK) 7 and GRK1 in mammalian cone photoreceptor cells: implications for cone cell phototransduction. *J. Neurosci.*, **21**, 9175-9184 (2001)
- [190] Sturany, S.; Van Lint, J.; Muller, F.; Wilda, M.; Hameister, H.; Hocker, M.; Brey, A.; Gern, U.; Vandenheede, J.; Gress, T.; Adler, G.; Seufferlein, T.: Molecular cloning and characterization of the human protein kinase D2. A novel member of the protein kinase D family of serine threonine kinases. *J. Biol. Chem.*, **276**, 3310-3318 (2001)
- [191] Sakurai, Y.; Onishi, Y.; Tanimoto, Y.; Kizaki, H.: Novel protein kinase C δ isoform insensitive to caspase-3. *Biol. Pharm. Bull.*, **24**, 973-977 (2001)
- [192] Matsubara, A.; Wasson, J.C.; Donelan, S.S.; Welling, C.M.; Glaser, B.; Permutt, M.A.: Isolation and characterization of the human AKT1 gene, identification of 13 single nucleotide polymorphisms (SNPs), and their lack of association with Type II diabetes. *Diabetologia*, **44**, 910-913 (2001)
- [198] Sutton, R.B.; Sprang, S.R.: Structure of the protein kinase C β phospholipid-binding C₂ domain complexed with Ca²⁺. *Structure*, **6**, 1395-1405 (1998)
- [199] Keranen, L.M.; Dutil, E.M.; Newton, A.C.: Protein kinase C is regulated in vivo by three functionally distinct phosphorylations. *Curr. Biol.*, **5**, 1394-1403 (1995)
- [200] Zhang, J.; Wang, L.; Schwartz, J.; Bond, R.W.; Bishop, W.R.: Phosphorylation of Thr642 is an early event in the processing of newly synthesized protein kinase C β 1 and is essential for its activation. *J. Biol. Chem.*, **269**, 19578-19584 (1994)

- [201] Orr, J.W.; Newton, A.C.: Requirement for negative charge on "activation loop" of protein kinase C. *J. Biol. Chem.*, **269**, 27715-27718 (1994)
- [202] Zhang, J.; Wang, L.; Petrin, J.; Bishop, W.R.; Bond, R.W.: Characterization of site-specific mutants altered at protein kinase C β 1 isozyme autophosphorylation sites. *Proc. Natl. Acad. Sci. USA*, **90**, 6130-6134 (1993)
- [203] Tang, Y.M.; Ashendel, C.L.: Isolation of cloned mouse protein kinase C β -II cDNA and its sequence. *Nucleic Acids Res.*, **18**, 5310 (1990)
- [204] Housey, G.M.; Johnson, M.D.; Hsiao, W.L.; O'Brian, C.A.; Murphy, J.P.; Kirschmeier, P.; Weinstein, I.B.: Overproduction of protein kinase C causes disordered growth control in rat fibroblasts. *Cell*, **52**, 343-354 (1988)
- [205] Housey, G.M.; O'Brian, C.A.; Johnson, M.D.; Kirschmeier, P.; Weinstein, I.B.: Isolation of cDNA clones encoding protein kinase C: evidence for a protein kinase C-related gene family. *Proc. Natl. Acad. Sci. USA*, **84**, 1065-1069 (1987)
- [206] Ono, Y.; Kurokawa, T.; Kawahara, K.; Nishimura, O.; Marumoto, R.; Igarashi, K.; Sugino, Y.; Kikkawa, U.; Ogita, K.; Nishizuka, Y.: Cloning of rat brain protein kinase C complementary DNA. *FEBS Lett.*, **203**, 111-115 (1986)
- [207] Ono, Y.; Kurokawa, T.; Fujii, T.; Kawahara, K.; Igarashi, K.; Kikkawa, U.; Ogita, K.; Nishizuka, Y.: Two types of complementary DNAs of rat brain protein kinase C. Heterogeneity determined by alternative splicing. *FEBS Lett.*, **206**, 347-352 (1986)
- [208] Knopf, J.L.; Lee, M.H.; Sultzman, L.A.; Kriz, R.W.; Loomis, C.R.; Hewick, R.M.; Bell, R.M.: Cloning and expression of multiple protein kinase C cDNAs. *Cell*, **46**, 491-502 (1986)
- [209] Le Good, J.A.; Ziegler, W.H.; Parekh, D.B.; Alessi, D.R.; Cohen, P.; Parker, P.J.: Protein kinase C isotypes controlled by phosphoinositide 3-kinase through the protein kinase PDK1. *Science*, **281**, 2042-2045 (1998)
- [210] Zhang, G.; Kazanietz, M.G.; Blumberg, P.M.; Hurley, J.H.: Crystal structure of the cys2 activator-binding domain of protein kinase C δ in complex with phorbol ester. *Cell*, **81**, 917-924 (1995)
- [211] Mizuno, K.; Kubo, K.; Saido, T.C.; Akita, Y.; Osada, S.; Kuroki, T.; Ohno, S.; Suzuki, K.: Structure and properties of a ubiquitously expressed protein kinase C, nPKC δ . *Eur. J. Biochem.*, **202**, 931-940 (1991)
- [212] Mischak, H.; Bodenteich, A.; Kolch, W.; Goodnight, J.; Hofer, F.; Mushinski, J.F.: Mouse protein kinase C- δ , the major isoform expressed in mouse hemopoietic cells: sequence of the cDNA, expression patterns, and characterization of the protein. *Biochemistry*, **30**, 7925-7931 (1991)
- [213] Brunet, A.; Park, J.; Tran, H.; Hu, L.S.; Hemmings, B.A.; Greenberg, M.E.: Protein kinase SGK mediates survival signals by phosphorylating the forkhead transcription factor FKHRL1 (FOXO3a). *Mol. Cell. Biol.*, **21**, 952-965 (2001)
- [214] Lang, F.; Klingel, K.; Wagner, C.A.; et al: Deranged transcriptional regulation of cell-volume-sensitive kinase hSGK in diabetic nephropathy. *Proc. Natl. Acad. Sci. USA*, **97**, 8157-8162 (2000)

- [215] Kobayashi, T.; Cohen, P.: Activation of serum- and glucocorticoid-regulated protein kinase by agonists that activate phosphatidylinositol 3-kinase is mediated by 3-phosphoinositide-dependent protein kinase-1 (PDK1) and PDK2. *Biochem. J.*, **339**, 319-328 (1999)
- [216] Kobayashi, T.; Deak, M.; Morrice, N.; Cohen, P.: Characterization of the structure and regulation of two novel isoforms of serum- and glucocorticoid-induced protein kinase. *Biochem. J.*, **344**, 189-197 (1999)
- [217] Waldegger, S.; Erdel, M.; Nagl, U.O.; Barth, P.; Raber, G.; Steuer, S.; Utermann, G.; Paulmichl, M.; Lang, F.: Genomic organization and chromosomal localization of the human SGK protein kinase gene. *Genomics*, **51**, 299-302 (1998)
- [218] Waldegger, S.; Barth, P.; Raber, G.; Lang, F.: Cloning and characterization of a putative human serine/threonine protein kinase transcriptionally modified during anisotonic and isotonic alterations of cell volume. *Proc. Natl. Acad. Sci. USA*, **94**, 4440-4445 (1997)
- [220] Delcommenne, M.; Tan, C.; Gray, V.; Rue, L.; Woodgett, J.; Dedhar, S.: Phosphoinositide-3-OH kinase-dependent regulation of glycogen synthase kinase 3 and protein kinase B/AKT by the integrin-linked kinase. *Proc. Natl. Acad. Sci. USA*, **95**, 11211-11216 (1998)
- [221] Alessi, D.R.; Andjelkovic, M.; Caudwell, B.; Cron, P.; Morrice, N.; Cohen, P.; Hemmings, B.A.: Mechanism of activation of protein kinase B by insulin and IGF¹. *EMBO J.*, **15**, 6541-6551 (1996)
- [222] Coffey, P.J.; Woodgett, J.R.: Molecular cloning and characterisation of a novel putative protein-serine kinase related to the cAMP-dependent and protein kinase C families. *Eur. J. Biochem.*, **205**, 1217 (1992)
- [223] Jones, P.F.; Jakubowicz, T.; Pitossi, F.J.; Maurer, F.; Hemmings, B.A.: Molecular cloning and identification of a serine/threonine protein kinase of the second-messenger subfamily. *Proc. Natl. Acad. Sci. USA*, **88**, 4171-4175 (1991)
- [224] Coffey, P.J.; Woodgett, J.R.: Molecular cloning and characterisation of a novel putative protein-serine kinase related to the cAMP-dependent and protein kinase C families. *Eur. J. Biochem.*, **201**, 475-481 (1991)
- [225] Brodbeck, D.; Hill, M.M.; Hemmings, B.A.: Two splice variants of protein kinase B γ have different regulatory capacity depending on the presence or absence of the regulatory phosphorylation site serine 472 in the carboxyl-terminal hydrophobic domain. *J. Biol. Chem.*, **276**, 29550-29558 (2001)
- [226] Nakatani, K.; Sakaue, H.; Thompson, D.A.; Weigel, R.J.; Roth, R.A.: Identification of a human Akt3 (protein kinase B γ) which contains the regulatory serine phosphorylation site. *Biochem. Biophys. Res. Commun.*, **257**, 906-910 (1999)
- [227] Masure, S.; Haefner, B.; Wesselink, J.J.; Hoefnagel, E.; Mortier, E.; Verhaselt, P.; Tuytelaars, A.; Gordon, R.; Richardson, A.: Molecular cloning, expression and characterization of the human serine/threonine kinase Akt-3. *Eur. J. Biochem.*, **265**, 353-360 (1999)

- [228] Brodbeck, D.; Cron, P.; Hemmings, B.A.: A human protein kinase By with regulatory phosphorylation sites in the activation loop and in the C-terminal hydrophobic domain. *J. Biol. Chem.*, **274**, 9133-9136 (1999)
- [229] Zhang, Q.H.; Ye, M.; Wu, X.Y.; et al.: Cloning and functional analysis of cDNAs with open reading frames for 300 previously undefined genes expressed in CD34+ hematopoietic stem/progenitor cells. *Genome Res.*, **10**, 1546-1560 (2000)
- [241] Hao, C.M.; Breyer, R.M.; Davis, L.S.; Breyer, M.D.: Intrarenal distribution of rabbit PKC ζ . *Kidney Int.*, **51**, 1831-1837 (1997)
- [242] Kothe, G.O.; Free, S.J.: The isolation and characterization of nrc-1 and nrc-2, two genes encoding protein kinases that control growth and development in *Neurospora crassa*. *Genetics*, **149**, 117-130 (1998)
- [243] Salanoubat, M.; Lemcke, K.; Rieger, M.; Ansorge, W.; Unseld, M.; et al.: Sequence and analysis of chromosome 3 of the plant *Arabidopsis thaliana*. *Nature*, **408**, 820-822 (2000)
- [244] Christie, J.M.; Salomon, M.; Nozue, K.; Wada, M.; Briggs, W.R.: LOV (light, oxygen, or voltage) domains of the blue-light photoreceptor phototropin (nph1): binding sites for the chromophore flavin mononucleotide. *Proc. Natl. Acad. Sci. USA*, **96**, 8779-8783 (1999)
- [245] Christie, J.M.; Reymond, P.; Powell, G.K.; Bernasconi, P.; Raibekas, A.A.; Liscum, E.; Briggs, W.R.: *Arabidopsis* NPH1: a flavoprotein with the properties of a photoreceptor for phototropism. *Science*, **282**, 1698-1701 (1998)
- [246] Huala, E.; Oeller, P.W.; Liscum, E.; Han, I.S.; Larsen, E.; Briggs, W.R.: *Arabidopsis* NPH1: a protein kinase with a putative redox-sensing domain. *Science*, **278**, 2120-2123 (1997)
- [257] Hayashi, A.; Seki, N.; Hattori, A.; Kozuma, S.; Saito, T.: PKCnu, a new member of the protein kinase C family, composes a fourth subfamily with PKCmu. *Biochim. Biophys. Acta*, **1450**, 99-106 (1999)
- [263] Nishizuka, Y.: The molecular heterogeneity of protein kinase C and its implications for cellular regulation. *Nature*, **334**, 661-665 (1988)
- [264] Parker, P.J.; Coussens, L.; Totty, N.; Rhee, L.; Young, S.; Chen, E.; Stabel, S.; Waterfield, M.D.; Ullrich, A.: The complete primary structure of protein kinase C - the major phorbol ester receptor. *Science*, **233**, 853-859 (1986)
- [265] Coussens, L.; Parker, P.J.; Rhee, L.; Yang-Feng, T.L.; Chen, E.; Waterfield, M.D.; Francke, U.; Ullrich, A.: Multiple, distinct forms of bovine and human protein kinase C suggest diversity in cellular signaling pathways. *Science*, **233**, 859-866 (1986)
- [266] Dryja, T.P.; McEvoy, J.; McGee, T.L.; Berson, E.L.: No mutations in the coding region of the PRKCG gene in three families with retinitis pigmentosa linked to the RP11 locus on chromosome 19q. *Am. J. Hum. Genet.*, **65**, 926-928 (1999)
- [267] Al-Magthteh, M.; Vithana, E.N.; Inglehearn, C.F.; Moore, T.; Bird, A.C.; Bhattacharya, S.S.: Segregation of a PRKCG mutation in two RP11 families. *Am. J. Hum. Genet.*, **62**, 1248-1252 (1998)
- [268] Kochs, G.; Hummel, R.; Meyer, D.; Hug, H.; Marme, D.; Sarre, T.F.: Activation and substrate specificity of the human protein kinase C α and ζ isoenzymes. *Eur. J. Biochem.*, **216**, 597-606 (1993)

- [269] Rosenthal, A.; Rhee, L.; Yadegari, R.; Paro, R.; Ullrich, A.; Goeddel, D.V.: Structure and nucleotide sequence of a *Drosophila melanogaster* protein kinase C gene. *EMBO J.*, **6**, 433-441 (1987)
- [270] Ono, Y.; Fujii, T.; Igarashi, K.; Kikkawa, U.; Ogita, K.; Nishizuka, Y.: Nucleotide sequences of cDNAs for α and γ subspecies of rat brain protein kinase C. *Nucleic Acids Res.*, **16**, 5199-5200 (1988)
- [271] Kikkawa, U.; Ogita, K.; Ono, Y.; Asaoka, Y.; Shearman, M.S.; Fujii, T.; Ase, K.; Sekiguchi, K.; Igarashi, K.; Nishizuka, Y.: The common structure and activities of four subspecies of rat brain protein kinase C family. *FEBS Lett.*, **223**, 212-216 (1987)
- [272] Xu, R.X.; Pawelczyk, T.; Xia, T.H.; Brown, S.C.: NMR structure of a protein kinase C- γ phorbol-binding domain and study of protein-lipid micelle interactions. *Biochemistry*, **36**, 10709-10717 (1997)
- [273] Bowers, B.J.; Parham, C.L.; Sikela, J.M.; Wehner, J.M.: Isolation and sequence of a mouse brain cDNA coding for protein kinase C- γ isozyme. *Gene*, **123**, 263-265 (1993)
- [274] Chen, K.H.; Widen, S.G.; Wilson, S.H.; Huang, K.P.: Characterization of the 5'-flanking region of the rat protein kinase C γ gene. *J. Biol. Chem.*, **265**, 19961-19965 (1990)
- [275] Loftus, B.J.; Kim, U.J.; Sneddon, V.P.; Kalush, F.; Brandon, R.; Fuhrmann, J.; Mason, T.; Crosby, M.L.; Barnstead, M.; Cronin, L.; Deslattes Mays, A.; Cao, Y.; Xu, R.X.; Kang, H.L.; Mitchell, S.; Eichler, E.E.; Harris, P.C.; Venter, J.C.; Adams, M.D.: Genome duplications and other features in 12 Mb of DNA sequence from human chromosome 16p and 16q. *Genomics*, **60**, 295-308 (1999)
- [276] Mahajna, J.; King, P.; Parker, P.; Haley, J.: Autoregulation of cloned human protein kinase C β and γ gene promoters in U937 cells. *DNA Cell Biol.*, **14**, 213-222 (1995)
- [277] Obeid, L.M.; Blobe, G.C.; Karolak, L.A.; Hannun, Y.A.: Cloning and characterization of the major promoter of the human protein kinase C β gene. Regulation by phorbol esters. *J. Biol. Chem.*, **267**, 20804-20810 (1992)
- [278] Kubo, K.; Ohno, S.; Suzuki, K.: Nucleotide sequence of the 3' portion of a human gene for protein kinase C β I/ β II. *Nucleic Acids Res.*, **15**, 7179-7180 (1987)
- [279] Kubo, K.; Ohno, S.; Suzuki, K.: Primary structures of human protein kinase C β I and β II differ only in their C-terminal sequences. *FEBS Lett.*, **223**, 138-142 (1987)
- [280] Coussens, L.; Rhee, L.; Parker, P.J.; Ullrich, A.: Alternative splicing increases the diversity of the human protein kinase C family. *DNA*, **6**, 389-394 (1987)
- [281] Ohno, S.; Kawasaki, H.; Imajoh, S.; Suzuki, K.; Inagaki, M.; Yokokura, H.; Sakoh, T.; Hidaka, H.: Tissue-specific expression of three distinct types of rabbit protein kinase C. *Nature*, **325**, 161-166 (1987)
- [286] Ueyama, T.; Ren, Y.; Ohmori, S.; Sakai, K.; Tamaki, N.; Saito, N.: cDNA cloning of an alternative splicing variant of protein kinase C δ (PKC δ III), a new truncated form of PKC δ , in rats. *Biochem. Biophys. Res. Commun.*, **269**, 557-563 (2000)

- [287] Kurkinen, K.M.; Keinänen, R.A.; Karhu, R.; Koistinaho, J.: Genomic structure and chromosomal localization of the rat protein kinase C δ -gene. *Gene*, **242**, 115-123 (2000)
- [288] Pappa, H.; Murray-Rust, J.; Dekker, L.V.; Parker, P.J.; McDonald, N.Q.: Crystal structure of the C₂ domain from protein kinase C- δ . *Structure*, **6**, 885-894 (1998)
- [289] Garcia-Paramio, P.; Cabrerizo, Y.; Bornancin, F.; Parker, P.J.: The broad specificity of dominant inhibitory protein kinase C mutants infers a common step in phosphorylation. *Biochem. J.*, **333**, 631-636 (1998)
- [290] Ono, Y.; Fujii, T.; Ogita, K.; Kikkawa, U.; Igarashi, K.; Nishizuka, Y.: The structure, expression, and properties of additional members of the protein kinase C family. *J. Biol. Chem.*, **263**, 6927-6932 (1988)
- [291] Ono, Y.; Fujii, T.; Ogita, K.; Kikkawa, U.; Igarashi, K.; Nishizuka, Y.: Identification of three additional members of rat protein kinase C family: δ -, ϵ - and zeta-subspecies. *FEBS Lett.*, **226**, 125-128 (1987)
- [292] Ono, Y.; Fujii, T.; Ogita, K.; Kikkawa, U.; Igarashi, K.; Nishizuka, Y.: Protein kinase C zeta subspecies from rat brain: its structure, expression, and properties. *Proc. Natl. Acad. Sci. USA*, **86**, 3099-3103 (1989)
- [293] Jones, S.W.; Erikson, E.; Blenis, J.; Maller, J.L.; Erikson, R.L.: A *Xenopus* ribosomal protein S6 kinase has two apparent kinase domains that are each similar to distinct protein kinases. *Proc. Natl. Acad. Sci. USA*, **85**, 3377-3381 (1988)
- [294] Ohno, S.; Kawasaki, H.; Konno, Y.; Inagaki, M.; Hidaka, H.; Suzuki, K.: A fourth type of rabbit protein kinase C. *Biochemistry*, **27**, 2083-2087 (1988)
- [295] Ohno, S.; Akita, Y.; Konno, Y.; Imajoh, S.; Suzuki, K.: A novel phorbol ester receptor/protein kinase, nPKC, distantly related to the protein kinase C family. *Cell*, **53**, 731-741 (1988)
- [296] Chen, P.; Lee, K.S.; Levin, D.E.: A pair of putative protein kinase genes (YPK1 and YPK2) is required for cell growth in *Saccharomyces cerevisiae*. *Mol. Gen. Genet.*, **236**, 443-447 (1993)
- [297] Maurer, R.A.: Isolation of a yeast protein kinase gene by screening with a mammalian protein kinase cDNA. *DNA*, **7**, 469-474 (1988)
- [299] Schaeffer, E.; Smith, D.; Mardon, G.; Quinn, W.; Zuker, C.: Isolation and characterization of two new *Drosophila* protein kinase C genes, including one specifically expressed in photoreceptor cells. *Cell*, **57**, 403-412 (1989)
- [300] Lawton, M.A.; Yamamoto, R.T.; Hanks, S.K.; Lamb, C.J.: Molecular cloning of plant transcripts encoding protein kinase homologs. *Proc. Natl. Acad. Sci. USA*, **86**, 3140-3144 (1989)
- [301] Wang, Q.J.; Acs, P.; Goodnight, J.; Blumberg, P.M.; Mischak, H.; Mushinski, J.F.: The catalytic domain of PKC- ϵ , in reciprocal PKC- δ and - ϵ chimeras, is responsible for conferring tumorigenicity to NIH3T3 cells, whereas both regulatory and catalytic domains of PKC- ϵ contribute to in vitro transformation. *Oncogene*, **16**, 53-60 (1998)
- [302] Schaap, D.; Parker, P.J.; Bristol, A.; Kriz, R.; Knopf, J.: Unique substrate specificity and regulatory properties of PKC- ϵ : a rationale for diversity. *FEBS Lett.*, **243**, 351-357 (1989)

- [303] Kalderon, D.; Rubin, G.M.: Isolation and characterization of *Drosophila* cAMP-dependent protein kinase genes. *Genes Dev.*, **2**, 1539-1556 (1988)
- [304] McSwine-Kennick, R.L.; McKeegan, E.M.; Johnson, M.D.; Morin, M.J.: Phorbol diester-induced alterations in the expression of protein kinase C isozymes and their mRNAs. Analysis in wild-type and phorbol diester-resistant HL-60 cell clones. *J. Biol. Chem.*, **266**, 15135-15143 (1991)
- [305] Finkenzeller, G.; Marme, D.; Hug, H.: Sequence of human protein kinase C α . *Nucleic Acids Res.*, **18**, 2183 (1990)
- [306] Alcorta, D.A.; Crews, C.M.; Sweet, L.J.; Bankston, L.; Jones, S.W.; Erikson, R.L.: Sequence and expression of chicken and mouse rsk: homologs of *Xenopus laevis* ribosomal S6 kinase. *Mol. Cell. Biol.*, **9**, 3850-3859 (1989)
- [307] Kubo, K.; Ohno, S.; Matsumoto, S.; Yahara, I.; Suzuki, K.: A novel yeast gene coding for a putative protein kinase. *Gene*, **76**, 177-180 (1989)
- [308] Staudinger, J.; Zhou, J.; Burgess, R.; Elledge, S.J.; Olson, E.N.: PICK1: a perinuclear binding protein and substrate for protein kinase C isolated by the yeast two-hybrid system. *J. Cell Biol.*, **128**, 263-271 (1995)
- [309] Megidish, T.; Mazurek, N.: A mutant protein kinase C that can transform fibroblasts. *Nature*, **342**, 807-811 (1989)
- [310] Rose-John, S.; Dietrich, A.; Marks, F.: Molecular cloning of mouse protein kinase C (PKC) cDNA from Swiss 3T3 fibroblasts. *Gene*, **74**, 465-471 (1988)
- [311] Burnett, P.E.; Blackshaw, S.; Lai, M.M.; Qureshi, I.A.; Burnett, A.F.; Sabatini, D.M.; Snyder, S.H.: Neurabin is a synaptic protein linking p70 S6 kinase and the neuronal cytoskeleton. *Proc. Natl. Acad. Sci. USA*, **95**, 8351-8356 (1998)
- [312] Kozma, S.C.; Ferrari, S.; Bassand, P.; Siegmann, M.; Totty, N.; Thomas, G.: Cloning of the mitogen-activated S6 kinase from rat liver reveals an enzyme of the second messenger subfamily. *Proc. Natl. Acad. Sci. USA*, **87**, 7365-7369 (1990)
- [313] Harmann, B.; Kilimann, M.W.: cDNA encoding a 59 kDa homolog of ribosomal protein S6 kinase from rabbit liver. *FEBS Lett.*, **273**, 248-252 (1990)
- [314] Banerjee, P.; Ahmad, M.F.; Grove, J.R.; Kozlosky, C.; Price, D.J.; Avruch, J.: Molecular structure of a major insulin/mitogen-activated 70-kDa S6 protein kinase. *Proc. Natl. Acad. Sci. USA*, **87**, 8550-8554 (1990)
- [315] Beushausen, S.; Bayley, H.: A relative of the catalytic subunit of cyclic AMP-dependent protein kinase in *Aplysia* spermatozoa. *Mol. Cell. Biol.*, **10**, 6775-6780 (1990)
- [316] Toyn, J.H.; Araki, H.; Sugino, A.; Johnston, L.H.: The cell-cycle-regulated budding yeast gene DBF2, encoding a putative protein kinase, has a homologue that is not under cell-cycle control. *Gene*, **104**, 63-70 (1991)
- [317] Johnston, L.H.; Eberly, S.L.; Chapman, J.W.; Araki, H.; Sugino, A.: The product of the *Saccharomyces cerevisiae* cell cycle gene DBF2 has homology with protein kinases and is periodically expressed in the cell cycle. *Mol. Cell. Biol.*, **10**, 1358-1366 (1990)
- [326] Osada, S.; Mizuno, K.; Saido, T.C.; Akita, Y.; Suzuki, K.; Kuroki, T.; Ohno, S.: A phorbol ester receptor/protein kinase, nPKC ϵ , a new member of

- the protein kinase C family predominantly expressed in lung and skin. *J. Biol. Chem.*, **265**, 22434-22440 (1990)
- [327] Grove, J.R.; Banerjee, P.; Balasubramanyam, A.; Coffey, P.J.; Price, D.J.; Avruch, J.; Woodgett, J.R.: Cloning and expression of two human p70 S6 kinase polypeptides differing only at their amino termini. *Mol. Cell. Biol.*, **11**, 5541-5550 (1991)
- [331] Baymiller, J.; McCullough, J.E.: *Saccharomyces cerevisiae* cell lysis mutations *cly5* and *cly7* define temperature-sensitive alleles of PKC1, the gene encoding yeast protein kinase C. *Yeast*, **13**, 305-312 (1997)
- [332] Obermaier, B.; Gassenhuber, J.; Piravandi, E.; Domdey, H.: Sequence analysis of a 78.6 kb segment of the left end of *Saccharomyces cerevisiae* chromosome II. *Yeast*, **11**, 1103-1112 (1995)
- [333] Levin, D.E.; Fields, F.O.; Kunisawa, R.; Bishop, J.M.; Thorner, J.: A candidate protein kinase C gene, PKC1, is required for the *S. cerevisiae* cell cycle. *Cell*, **62**, 213-224 (1990)
- [334] Palmer, R.H.; Ridden, J.; Parker, P.J.: Identification of multiple, novel, protein kinase C-related gene products. *FEBS Lett.*, **356**, 5-8 (1994)
- [335] Bacher, N.; Zisman, Y.; Berent, E.; Livneh, E.: Isolation and characterization of PKC-L, a new member of the protein kinase C-related gene family specifically expressed in lung, skin, and heart. *Mol. Cell. Biol.*, **12**, 1404 (1992)
- [336] Bacher, N.; Zisman, Y.; Berent, E.; Livneh, E.: Isolation and characterization of PKC-L, a new member of the protein kinase C-related gene family specifically expressed in lung, skin, and heart. *Mol. Cell. Biol.*, **11**, 126-133 (1991)
- [337] Haribabu, B.; Dottin, R.P.: Identification of a protein kinase multigene family of *Dictyostelium discoideum*: molecular cloning and expression of a cDNA encoding a developmentally regulated protein kinase. *Proc. Natl. Acad. Sci. USA*, **88**, 1115-1119 (1991)
- [341] Bellacosa, A.; Franke, T.F.; Gonzalez-Portal, M.E.; Datta, K.; Taguchi, T.; Gardner, J.; Cheng, J.Q.; Testa, J.R.; Tsichlis, P.N.: Structure, expression and chromosomal mapping of *c-akt*: relationship to *v-akt* and its implications. *Oncogene*, **8**, 745-754 (1993)
- [342] Cheng, J.Q.; Godwin, A.K.; Bellacosa, A.; Taguchi, T.; Franke, T.F.; Hamilton, T.C.; Tsichlis, P.N.; Testa, J.R.: AKT2, a putative oncogene encoding a member of a subfamily of protein-serine/threonine kinases, is amplified in human ovarian carcinomas. *Proc. Natl. Acad. Sci. USA*, **89**, 9267-9271 (1992)
- [343] Jones, P.F.; Jakubowicz, T.; Hemmings, B.A.: Molecular cloning of a second form of *rac* protein kinase. *Cell Regul.*, **2**, 1001-1009 (1991)
- [344] Sallese, M.; Mariggio, S.; Collodel, G.; Moretti, E.; Piomboni, P.; Baccetti, B.; De Blasi, A.: G protein-coupled receptor kinase GRK4. Molecular analysis of the four isoforms and ultrastructural localization in spermatozoa and germinal cells. *J. Biol. Chem.*, **272**, 10188-10195 (1997)
- [345] Sallese, M.; Lombardi, M.S.; De Blasi, A.: Two isoforms of G protein-coupled receptor kinase 4 identified by molecular cloning. *Biochem. Biophys. Res. Commun.*, **199**, 848-854 (1994)

- [346] Ambrose, C.; James, M.; Barnes, G.; Lin, C.; Bates, G.; Altherr, M.; Duyao, M.; Groot, N.; Church, D.; Wasmuth, J.J.; et al.: A novel G protein-coupled receptor kinase gene cloned from 4p16.3. *Hum. Mol. Genet.*, **1**, 697-703 (1992)
- [348] Cassill, J.A.; Whitney, M.; Joazeiro, C.A.; Becker, A.; Zuker, C.S.: Isolation of *Drosophila* genes encoding G protein-coupled receptor kinases. *Proc. Natl. Acad. Sci. USA*, **88**, 11067-11070 (1991)
- [349] Schneider, L.E.; Spradling, A.C.: The *Drosophila* G-protein-coupled receptor kinase homologue Gprk2 is required for egg morphogenesis. *Development*, **124**, 2591-2602 (1997)
- [359] Sano, T.; Tabuse, Y.; Nishiwaki, K.; Miwa, J.: The tpa-1 gene of *Caenorhabditis elegans* encodes two proteins similar to Ca(2+)-independent protein kinase Cs: evidence by complete genomic and complementary DNA sequences of the tpa-1 gene. *J. Mol. Biol.*, **251**, 477-485 (1995)
- [360] Tabuse, Y.; Nishiwaki, K.; Miwa, J.: Mutations in a protein kinase C homolog confer phorbol ester resistance on *Caenorhabditis elegans*. *Science*, **243**, 1713-1716 (1989)
- [361] Land, M.; Islas-Trejo, A.; Freedman, J.H.; Rubin, C.S.: Structure and expression of a novel, neuronal protein kinase C (PKC1B) from *Caenorhabditis elegans*. PKC1B is expressed selectively in neurons that receive, transmit, and process environmental signals. *J. Biol. Chem.*, **269**, 9234-9244 (1994)
- [362] Kunapuli, P.; Benovic, J.L.: Cloning and expression of GRK5: a member of the G protein-coupled receptor kinase family. *Proc. Natl. Acad. Sci. USA*, **90**, 5588-5592 (1993)
- [364] Toda, T.; Shimanuki, M.; Yanagida, M.: Two novel protein kinase C-related genes of fission yeast are essential for cell viability and implicated in cell shape control. *EMBO J.*, **12**, 1987-1995 (1993)
- [365] Ding, D.Q.; Tomita, Y.; Yamamoto, A.; Chikashige, Y.; Haraguchi, T.; Hiraoaka, Y.: Large-scale screening of intracellular protein localization in living fission yeast cells by the use of a GFP-fusion genomic DNA library. *Genes Cells*, **5**, 169-190 (2000)
- [366] Mazzei, G.J.; Schmid, E.M.; Knowles, J.K.; Payton, M.A.; Maundrell, K.G.: A Ca(2+)-independent protein kinase C from fission yeast. *J. Biol. Chem.*, **268**, 7401-7406 (1993)
- [370] Yarden, O.; Plamann, M.; Ebbrole, D.J.; Yanofsky, C.: Cot-1, a gene required for hyphal elongation in *Neurospora crassa*, encodes a protein kinase. *EMBO J.*, **11**, 2159-2166 (1992)
- [376] Mazzearella, R.; Ciccodicola, A.; Esposito, T.; Arcucci, A.; Migliaccio, C.; Jones, C.; Schlessinger, D.; D'Urso, M.; D'Esposito, M.: Human protein kinase C iota gene (PRKCI) is closely linked to the BTK gene in Xq21.3. *Genomics*, **26**, 629-631 (1995)
- [377] Selbie, L.A.; Schmitz-Peiffer, C.; Sheng, Y.; Biden, T.J.: Molecular cloning and characterization of PKC iota, an atypical isoform of protein kinase C derived from insulin-secreting cells. *J. Biol. Chem.*, **268**, 24296-24302 (1993)

- [378] Mizoguchi, T.; Hayashida, N.; Yamaguchi-Shinozaki, K.; Kamada, H.; Shinozaki, K.: Two genes that encode ribosomal-protein S6 kinase homologs are induced by cold or salinity stress in *Arabidopsis thaliana*. *FEBS Lett.*, **358**, 199-204 (1995)
- [379] Zhang, S.H.; Broome, M.A.; Lawton, M.A.; Hunter, T.; Lamb, C.J.: Atpk1, a novel ribosomal protein kinase gene from *Arabidopsis*. II. Functional and biochemical analysis of the encoded protein. *J. Biol. Chem.*, **269**, 17593-17599 (1994)
- [380] Zhang, S.H.; Lawton, M.A.; Hunter, T.; Lamb, C.J.: Atpk1, a novel ribosomal protein kinase gene from *Arabidopsis*. I. Isolation, characterization, and expression. *J. Biol. Chem.*, **269**, 17586-17592 (1994)
- [381] Paravicini, G.; Mendoza, A.; Antonsson, B.; Cooper, M.; Losberger, C.; Payton, M.A.: The *Candida albicans* PKC1 gene encodes a protein kinase C homolog necessary for cellular integrity but not dimorphism. *Yeast*, **12**, 741-756 (1996)
- [382] Premont, R.T.; Koch, W.J.; Inglese, J.; Lefkowitz, R.J.: Identification, purification, and characterization of GRK5, a member of the family of G protein-coupled receptor kinases. *J. Biol. Chem.*, **269**, 6832-6841 (1994)
- [383] Stoffel, R.H.; Randall, R.R.; Premont, R.T.; Lefkowitz, R.J.; Inglese, J.: Palmitoylation of G protein-coupled receptor kinase, GRK6. Lipid modification diversity in the GRK family. *J. Biol. Chem.*, **269**, 27791-27794 (1994)
- [384] Benovic, J.L.; Gomez, J.: Molecular cloning and expression of GRK6. A new member of the G protein-coupled receptor kinase family. *J. Biol. Chem.*, **268**, 19521-19527 (1993)
- [390] Konishi, H.; Shinomura, T.; Kuroda, S.; Ono, Y.; Kikkawa, U.: Molecular cloning of rat RAC protein kinase α and β and their association with protein kinase C ζ . *Biochem. Biophys. Res. Commun.*, **205**, 817-825 (1994)
- [396] Abidi, F.; Jacquot, S.; Lassiter, C.; Trivier, E.; Hanauer, A.; Schwartz, C.E.: Novel mutations in Rsk-2, the gene for Coffin-Lowry syndrome (CLS). *Eur. J. Hum. Genet.*, **7**, 20-26 (1999)
- [397] Jacquot, S.; Merienne, K.; De Cesare, D.; Pannetier, S.; Mandel, J.L.; Sassone-Corsi, P.; Hanauer, A.: Mutation analysis of the RSK2 gene in Coffin-Lowry patients: extensive allelic heterogeneity and a high rate of de novo mutations. *Am. J. Hum. Genet.*, **63**, 1631-1640 (1998)
- [398] Trivier, E.; De Cesare, D.; Jacquot, S.; Pannetier, S.; Zackai, E.; Young, I.; Mandel, J.L.; Sassone-Corsi, P.; Hanauer, A.: Mutations in the kinase Rsk-2 associated with Coffin-Lowry syndrome. *Nature*, **384**, 567-570 (1996)
- [399] Bjorbaek, C.; Vik, T.A.; Echwald, S.M.; et al.: Cloning of a human insulin-stimulated protein kinase (ISPK-1) gene and analysis of coding regions and mRNA levels of the ISPK-1 and the protein phosphatase-1 genes in muscle from NIDDM patients. *Diabetes*, **44**, 90-97 (1995)
- [400] Moller, D.E.; Xia, C.H.; Tang, W.; Zhu, A.X.; Jakubowski, M.: Human rsk isoforms: cloning and characterization of tissue-specific expression. *Am. J. Physiol.*, **266**, C351-359 (1994)
- [401] Klink, A.; Schiebel, K.; Winkelmann, M.; et al.: The human protein kinase gene PKX1 on Xp22.3 displays Xp/Yp homology and is a site of chromosomal instability. *Hum. Mol. Genet.*, **4**, 869-878 (1995)

- [418] Groenen, P.J.; Wansink, D.G.; Coerwinkel, M.; van den Broek, W.; Jansen, G.; Wieringa, B.: Constitutive and regulated modes of splicing produce six major myotonic dystrophy protein kinase (DMPK) isoforms with distinct properties. *Hum. Mol. Genet.*, **9**, 605-616 (2000)
- [419] Mahadevan, M.S.; Amemiya, C.; Jansen, G.; Sabourin, L.; Baird, S.; Neville, C.E.; Wormskamp, N.; Segers, B.; Batzer, M.; Lamerdin, J.; et al.: Structure and genomic sequence of the myotonic dystrophy (DM kinase) gene. *Hum. Mol. Genet.*, **2**, 299-304 (1993)
- [420] Jansen, G.; Bartolomei, M.; Kalscheuer, V.; Merckx, G.; Wormskamp, N.; Mariman, E.; Smeets, D.; Ropers, H.H.; Wieringa, B.: No imprinting involved in the expression of DM-kinase mRNAs in mouse and human tissues. *Hum. Mol. Genet.*, **2**, 1221-1227 (1993)
- [421] Jansen, G.; Mahadevan, M.; Amemiya, C.; Wormskamp, N.; Segers, B.; Hendriks, W.; O'Hoy, K.; Baird, S.; Sabourin, L.; Lennon, G.; et al.: Characterization of the myotonic dystrophy region predicts multiple protein isoform-encoding mRNAs. *Nat. Genet.*, **1**, 261-266 (1992)
- [428] Fehr, C.; Fickova, M.; Hiemke, C.; Reuss, S.; Dahmen, N.: Molecular cloning of rat G-protein-coupled receptor kinase 6 (GRK6) from brain tissue, and its mRNA expression in different brain regions and peripheral tissues. *Brain Res. Mol. Brain Res.*, **49**, 278-282 (1997)
- [429] Morawetz, R.; Lendenfeld, T.; Mischak, H.; Muhlbauer, M.; Gruber, F.; Goodnight, J.; de Graaff, L.H.; Visser, J.; Mushinski, J.F.; Kubicek, C.P.: Cloning and characterisation of genes (pkc1 and pkcA) encoding protein kinase C homologues from *Trichoderma reesei* and *Aspergillus niger*. *Mol. Gen. Genet.*, **250**, 17-28 (1996)
- [434] Basta, P.; Strickland, M.B.; Holmes, W.; Loomis, C.R.; Ballas, L.M.; Burns, D.J.: Sequence and expression of human protein kinase C- ϵ . *Biochim. Biophys. Acta*, **1132**, 154-160 (1992)
- [436] Goodnight, J.; Kazanietz, M.G.; Blumberg, P.M.; Mushinski, J.F.; Mischak, H.: The cDNA sequence, expression pattern and protein characteristics of mouse protein kinase C-zeta. *Gene*, **122**, 305-311 (1992)
- [437] Chang, J.D.; Xu, Y.; Raychowdhury, M.K.; Ware, J.A.: Molecular cloning and expression of a cDNA encoding a novel isoenzyme of protein kinase C (nPKC). A new member of the nPKC family expressed in skeletal muscle, megakaryoblastic cells, and platelets. *J. Biol. Chem.*, **268**, 14208-14214 (1993)
- [438] Baier, G.; Telford, D.; Giampa, L.; Coggeshall, K.M.; Baier-Bitterlich, G.; Isakov, N.; Altman, A.: Molecular cloning and characterization of PKC theta, a novel member of the protein kinase C (PKC) gene family expressed predominantly in hematopoietic cells. *J. Biol. Chem.*, **268**, 4997-5004 (1993)
- [439] Barbee, J.L.; Deutscher, S.L.; Loomis, C.R.; Burns, D.J.: The cDNA sequence encoding human protein kinase C- ζ . *Gene*, **132**, 305-306 (1993)
- [440] Nomoto, S.; Watanabe, Y.; Ninomiya-Tsuji, J.; Yang, L.X.; Nagai, Y.; Kiuchi, K.; Hagiwara, M.; Hidaka, H.; Matsumoto, K.; Irie, K.: Functional analyses of mammalian protein kinase C isozymes in budding yeast and mammalian fibroblasts. *Genes Cells*, **2**, 601-614 (1997)

- [441] Aris, J.P.; Basta, P.V.; Holmes, W.D.; Ballas, L.M.; Moomaw, C.; Rankl, N.B.; Blobel, G.; Loomis, C.R.; Burns, D.J.: Molecular and biochemical characterization of a recombinant human PKC- δ family member. *Biochim. Biophys. Acta*, **1174**, 171-181 (1993)
- [442] Park, J.; Leong, M.L.; Buse, P.; Maiyar, A.C.; Firestone, G.L.; Hemmings, B.A.: Serum and glucocorticoid-inducible kinase (SGK) is a target of the PI 3-kinase-stimulated signaling pathway. *EMBO J.*, **18**, 3024-3033 (1999)
- [443] Richards, J.S.; Fitzpatrick, S.L.; Clemens, J.W.; Morris, J.K.; Alliston, T.; Sirois, J.: Ovarian cell differentiation: a cascade of multiple hormones, cellular signals, and regulated genes. *Recent Prog. Horm. Res.*, **50**, 223-254 (1995)
- [444] Imaizumi, K.; Tsuda, M.; Wanaka, A.; Tohyama, M.; Takagi, T.: Differential expression of *sgk* mRNA, a member of the Ser/Thr protein kinase gene family, in rat brain after CNS injury. *Brain Res. Mol. Brain Res.*, **26**, 189-196 (1994)
- [445] Webster, M.K.; Goya, L.; Ge, Y.; Maiyar, A.C.; Firestone, G.L.: Characterization of *sgk*, a novel member of the serine/threonine protein kinase gene family which is transcriptionally induced by glucocorticoids and serum. *Mol. Cell. Biol.*, **13**, 2031-2040 (1993)
- [449] Gennarelli, M.; Lucarelli, M.; Zelano, G.; Pizzuti, A.; Novelli, G.; Dallapiccola, B.: Different expression of the myotonin protein kinase gene in discrete areas of human brain. *Biochem. Biophys. Res. Commun.*, **216**, 489-494 (1995)
- [450] Sasagawa, N.; Sorimachi, H.; Maruyama, K.; Arahata, K.; Ishiura, S.; Suzuki, K.: Expression of a novel human myotonin protein kinase (MtPK) cDNA clone which encodes a protein with a thymopoietin-like domain in COS cells. *FEBS Lett.*, **351**, 22-26 (1994)
- [451] Shaw, D.J.; McCurrach, M.; Rundle, S.A.; Harley, H.G.; Crow, S.R.; Sohn, R.; Thirion, J.P.; Hamshere, M.G.; Buckler, A.J.; Harper, P.S.; et al.: Genomic organization and transcriptional units at the myotonic dystrophy locus. *Genomics*, **18**, 673-679 (1993)
- [452] Fu, Y.H.; Friedman, D.L.; Richards, S.; Pearlman, J.A.; Gibbs, R.A.; Pizzuti, A.; Ashizawa, T.; Perryman, M.B.; Scarlato, G.; Fenwick, R.G., Jr.; et al.: Decreased expression of myotonin-protein kinase messenger RNA and protein in adult form of myotonic dystrophy. *Science*, **260**, 235-238 (1993)
- [453] Fu, Y.H.; Pizzuti, A.; Fenwick, R.G., Jr.; King, J.; Rajnarayan, S.; Dunne, P.W.; Dubel, J.; Nasser, G.A.; Ashizawa, T.; de Jong, P.; et al.: An unstable triplet repeat in a gene related to myotonic muscular dystrophy. *Science*, **255**, 1256-1258 (1992)
- [454] Brook, J.D.; McCurrach, M.E.; Harley, H.G.; Buckler, A.J.; Church, D.; Aburatani, H.; Hunter, K.; Stanton, V.P.; Thirion, J.P.; Hudson, T.; et al.: Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell*, **68**, 799-808 (1992)
- [462] Johannes, F.J.; Prestle, J.; Eis, S.; Oberhagemann, P.; Pfizenmaier, K.: PKC ϵ is a novel, atypical member of the protein kinase C family. *J. Biol. Chem.*, **269**, 6140-6148 (1994)

- [463] Zhao, Y.; Bjorbaek, C.; Weremowicz, S.; Morton, C.C.; Moller, D.E.: RSK3 encodes a novel pp90rsk isoform with a unique N-terminal sequence: growth factor-stimulated kinase function and nuclear translocation. *Mol. Cell. Biol.*, **15**, 4353-4363 (1995)
- [469] Palmer, R.H.; Ridden, J.; Parker, P.J.: Cloning and expression patterns of two members of a novel protein-kinase-C-related kinase family. *Eur. J. Biochem.*, **227**, 344-351 (1995)
- [470] Mukai, H.; Ono, Y.: A novel protein kinase with leucine zipper-like sequences: its catalytic domain is highly homologous to that of protein kinase C. *Biochem. Biophys. Res. Commun.*, **199**, 897-904 (1994)
- [471] Pepio, A.M.; Fan, X.; Sossin, W.S.: The role of C₂ domains in Ca²⁺-activated and Ca²⁺-independent protein kinase Cs in aplysia. *J. Biol. Chem.*, **273**, 19040-19048 (1998)
- [472] Sossin, W.S.; Diaz-Arrastia, R.; Schwartz, J.H.: Characterization of two isoforms of protein kinase C in the nervous system of *Aplysia californica*. *J. Biol. Chem.*, **268**, 5763-5768 (1993)
- [473] Kruger, K.E.; Sossin, W.S.; Sacktor, T.C.; Bergold, P.J.; Beushausen, S.; Schwartz, J.H.: Cloning and characterization of Ca²⁺-dependent and Ca²⁺-independent PKCs expressed in *Aplysia* sensory cells. *J. Neurosci.*, **11**, 2303-2313 (1991)
- [474] Pepio, A.M.; Sossin, W.S.: The C₂ domain of the Ca²⁺-independent protein kinase C Apl II inhibits phorbol ester binding to the C₁ domain in a phosphatidic acid-sensitive manner. *Biochemistry*, **37**, 1256-1263 (1998)
- [475] Sossin, W.S.; Schwartz, J.H.: Ca²⁺-independent protein kinase Cs contain an amino-terminal domain similar to the C₂ consensus sequence. *Trends Biochem. Sci.*, **18**, 207-208 (1993)
- [480] Altomare, D.A.; Guo, K.; Cheng, J.Q.; Sonoda, G.; Walsh, K.; Testa, J.R.: Cloning, chromosomal localization and expression analysis of the mouse Akt2 oncogene. *Oncogene*, **11**, 1055-1060 (1995)
- [481] Akimoto, K.; Mizuno, K.; Osada, S.; Hirai, S.; Tanuma, S.; Suzuki, K.; Ohno, S.: A new member of the third class in the protein kinase C family, PKC lambda, expressed dominantly in an undifferentiated mouse embryonal carcinoma cell line and also in many tissues and cells. *J. Biol. Chem.*, **269**, 12677-12683 (1994)
- [482] Matthews, S.A.; Rozengurt, E.; Cantrell, D.: Characterization of serine 916 as an in vivo autophosphorylation site for protein kinase D/Protein kinase Cmu. *J. Biol. Chem.*, **274**, 26543-26549 (1999)
- [483] Valverde, A.M.; Sinnett-Smith, J.; Van Lint, J.; Rozengurt, E.: Molecular cloning and characterization of protein kinase D: a target for diacylglycerol and phorbol esters with a distinctive catalytic domain. *Proc. Natl. Acad. Sci. USA*, **91**, 8572-8576 (1994)
- [484] Nagayama, Y.; Tanaka, K.; Hara, T.; Namba, H.; Yamashita, S.; Taniyama, K.; Niwa, M.: Involvement of G protein-coupled receptor kinase 5 in homologous desensitization of the thyrotropin receptor. *J. Biol. Chem.*, **271**, 10143-10148 (1996)

- [486] Morrice, N.A.; Gabrielli, B.; Kemp, B.E.; Wettenhall, R.E.: A cardiolipin-activated protein kinase from rat liver structurally distinct from the protein kinases C. *J. Biol. Chem.*, **269**, 20040-20046 (1994)
- [487] Konishi, H.; Kuroda, S.; Tanaka, M.; Matsuzaki, H.; Ono, Y.; Kameyama, K.; Haga, T.; Kikkawa, U.: Molecular cloning and characterization of a new member of the RAC protein kinase family: association of the pleckstrin homology domain of three types of RAC protein kinase with protein kinase C subspecies and β γ subunits of G proteins. *Biochem. Biophys. Res. Commun.*, **216**, 526-534 (1995)
- [488] Grove, J.R.; Price, D.J.; Banerjee, P.; Balasubramanyam, A.; Ahmad, M.F.; Avruch, J.: Regulation of an epitope-tagged recombinant Rsk-1 S6 kinase by phorbol ester and erk/MAP kinase. *Biochemistry*, **32**, 7727-7738 (1993)
- [489] Dekker, L.V.; Parker, P.J.; McIntyre, P.: Biochemical properties of rat protein kinase C- η expressed in COS cells. *FEBS Lett.*, **312**, 195-199 (1992)
- [497] Lee-Fruman, K.K.; Kuo, C.J.; Lippincott, J.; Terada, N.; Blenis, J.: Characterization of S6K2, a novel kinase homologous to S6K1. *Oncogene*, **18**, 5108-5114 (1999)
- [498] Saitoh, M.; ten Dijke, P.; Miyazono, K.; Ichijo, H.: Cloning and characterization of p70(S6K β) defines a novel family of p70 S6 kinases. *Biochem. Biophys. Res. Commun.*, **253**, 470-476 (1998)
- [499] Gout, I.; Minami, T.; Hara, K.; Tsujishita, Y.; Filonenko, V.; Waterfield, M.D.; Yonezawa, K.: Molecular cloning and characterization of a novel p70 S6 kinase, p70 S6 kinase β containing a proline-rich region. *J. Biol. Chem.*, **273**, 30061-30064 (1998)
- [500] Yntema, H.G.; van den Helm, B.; Kissing, J.; et al.: A novel ribosomal S6-kinase (RSK4; RPS6KA6) is commonly deleted in patients with complex X-linked mental retardation. *Genomics*, **62**, 332-343 (1999)
- [501] Kispert, A.; Stoger, R.J.; Caparros, M.; Herrmann, B.G.: The mouse Rsk3 gene maps to the Lehm66 elements carrying the t-complex responder Tcr. *Mamm. Genome*, **10**, 794-802 (1999)
- [504] Shima, H.; Pende, M.; Chen, Y.; Fumagalli, S.; Thomas, G.; Kozma, S.C.: Disruption of the p70(s6k)/p85(s6k) gene reveals a small mouse phenotype and a new functional S6 kinase. *EMBO J.*, **17**, 6649-6659 (1998)